

Singular Value Approximation and Reducing Directed to Undirected Graph Sparsification

AmirMahdi Ahmadinejad Amazon* ahmadi@alumni.stanford.edu	John Peebles Apple peebles@apple.com	Edward Pyne [†] MIT epyne@mit.edu
Aaron Sidford Stanford sidford@stanford.edu	Salil Vadhan [‡] Harvard University salil_vadhan@harvard.edu	

February 1, 2023

Abstract

In this paper, we introduce a new, spectral notion of approximation between directed graphs, which we call *Singular Value (SV) approximation*. SV-approximation is stronger than previous notions of spectral approximation considered in the literature, including spectral approximation of Laplacians for undirected graphs [ST04], standard approximation for directed graphs [CKP⁺17], and unit-circle approximation for directed graphs [AKM⁺20]. Moreover, SV approximation enjoys several useful properties not known to be possessed by previous notions of approximation, such as being preserved under products of random-walk matrices and with matrices of bounded norm.

Notably, we show that there is a simple black-box reduction from SV-sparsifying Eulerian directed graphs to SV-sparsifying undirected graphs. With this reduction in hand, we provide a nearly linear-time algorithm for SV-sparsifying undirected and hence also Eulerian directed graphs. This also yields the first nearly linear-time algorithm for unit-circle-sparsifying Eulerian directed graphs. In addition, we give a nearly linear-time algorithm for SV-sparsifying (and UC-sparsifying) random-walk polynomials of Eulerian directed graphs with second normalized singular value bounded away from 1 by $1/\text{poly}(n)$.

Finally, we show that a simple repeated-squaring and sparsification algorithm for solving Laplacian systems, introduced by Peng and Spielman [PS14] for undirected graphs, also works for Eulerian digraphs whose random-walk matrix is normal (i.e. unitarily diagonalizable), if we use SV-sparsification at each step. Prior Laplacian solvers for Eulerian digraphs are significantly more complicated.

*The work was started prior to joining Amazon and does not relate to Amazon

[†]Supported by an Akamai Presidential Fellowship.

[‡]Supported by a Simons Investigator Award.

Contents

1	Introduction	3
1.1	Our Contribution: Singular-Value Approximation	4
1.2	Comparison to Previous Notions of Approximation	5
1.3	Properties of SV Approximation	8
1.4	Algorithmic Results	10
1.5	Open Problems	12
1.6	Roadmap	12
2	Preliminaries	13
3	Singular Value Approximation	13
3.1	Matrix Approximation	13
3.2	Equivalent definitions of SV approximation	15
3.3	Properties of Singular Value Approximation	17
4	Sparsification	19
4.1	Cycle and Expander Decompositions	21
4.2	Sparsifying Bipartite Expanders	22
4.3	Sparsifying a Constant Fraction of Edges	24
4.4	Nearly Linear-Sized Sparsifiers	25
4.5	Derandomized Square Sparsification	26
4.6	Sparsification of Random Walk Polynomials	27
5	Squaring algorithm for solving normal directed Laplacians	31
A	SV Approximation Proofs	39
A.1	Equivalences	40
A.2	Properties	44
A.3	Separations	48
B	Facts about Singular Values	49

1 Introduction

Given an undirected graph, its associated *Laplacian* matrix is $\mathbf{L} = \mathbf{D} - \mathbf{A}$ where \mathbf{D} is the diagonal matrix of degrees of the graph and \mathbf{A} is its associated adjacency matrix. The linear algebraic properties of Laplacian matrices capture fundamental information about the corresponding graph, including random walks, cuts, and more. Correspondingly, starting with the seminal work of Spielman and Teng [ST04], nearly-linear time algorithms for solving Laplacian linear systems and related problems [ST04, BSS12, CKP⁺16, CKP⁺17, CKK⁺18] have been established as an important tool in algorithmic graph theory. These results have enabled asymptotic improvements for a variety of problems, such as max-flow and min-cut [CKM⁺11, LRS13, KLOS14, vdBGJ⁺22, vdBLL⁺21], traveling salesman [AGM⁺10, AG15], sampling random spanning trees [KM09, MST15], graph sparsification [SS08, LS13, KMP14, CKM⁺14, LS18, ABKS21, JS21], multicommodity flow [KMP12, KLOS14], and solving clustering and partitioning problems [AM85, KVV04, ACL06, OSV12].

This “Laplacian paradigm” was initially formulated for undirected graphs. However, there is a natural generalization of the Laplacian matrix to directed graphs, which analogously captures the information about random walks on the directed graph [CKP⁺16]. This generalization is $\mathbf{L} = \mathbf{D}_{\text{out}} - \mathbf{A}^\top$ where \mathbf{D}_{out} is now the diagonal matrix of out-degrees and \mathbf{A}^\top is the transpose of the adjacency matrix \mathbf{A} . It is called the *directed* Laplacian matrix, to emphasize that the graph corresponding to it is directed.

While the generalization of the *definition* of the Laplacian matrix to directed graphs is straightforward, the generalization of *algorithms* for Laplacian matrices is not so straightforward. Nonetheless, a sequence of papers [CKP⁺16, CKP⁺17, CKK⁺18] obtained nearly linear time algorithms for sparsifying and solving directed Laplacian linear systems. In order to do so, they needed to overcome three main difficulties not present in the undirected setting:

Challenge #1: While the kernel of an undirected Laplacian matrix is the all ones vector, computing the kernel of a directed Laplacian matrix \mathbf{L} corresponds to computing the stationary distribution π of the random walk on the directed graph ($\mathbf{L}\mathbf{D}_{\text{out}}^{-1}\pi = 0$). Without explicitly knowing the kernel, it is not known how to efficiently perform any kind of useful sparsification or approximately solve systems in \mathbf{L} . This difficulty was overcome in [CKP⁺16, AJSS19] which provide reductions from solving general directed Laplacian systems to the case where the graph is *Eulerian*, meaning that every vertex has the same in-degree as out-degree. In Eulerian graphs, the stationary distribution is simply proportional to the vertex degrees and the all ones vector is both the left and right kernels of the associated directed Laplacian.

Challenge #2: Undirected Laplacians \mathbf{L} are not only symmetric but also positive semidefinite (PSD), i.e. $x^\top \mathbf{L}x \geq 0$ for all x . This leads to a natural definition of multiplicative approximation between such matrices, namely saying that $\tilde{\mathbf{L}}$ is an ε -approximation of \mathbf{L} if $(1 - \varepsilon)\mathbf{L} \preceq \tilde{\mathbf{L}} \preceq (1 + \varepsilon)\mathbf{L}$, where \preceq is the Löwner order on PSD matrices. However, even though Laplacians of directed graphs are potentially asymmetric, the quadratic form $x^\top \mathbf{L}x$ depends only on a symmetrization of the Laplacian ($x^\top \mathbf{L}x = x^\top (1/2)(\mathbf{L} + \mathbf{L}^\top)x$). Consequently, the quadratic form discards key information about the associated directed graph (e.g. the quadratic form of a directed cycle and an undirected cycle are the same). Thus, defining approximation for directed graphs (even Eulerian ones) is more challenging than for undirected graphs and more complex notions of approximations were introduced in [CKP⁺16, AKM⁺20]. This additional complexity suggests that one formulate new sparsification algorithms that take into account the directedness of the graph.

Challenge #3: Finally, the recursive identities used for solving undirected Laplacian systems, while behaving nicely with respect to PSD approximation, do not behave as nicely with respect to the new directed matrix approximation definitions mentioned in Challenge #2. This necessitates different, more sophisticated recursion with a more involved analysis of the error [CKP⁺17, CKK⁺18, AKM⁺20, KMG22].

1.1 Our Contribution: Singular-Value Approximation

In this paper, we present a stronger and more robust notion for addressing Challenge #2 and use it to make progress in simplifying approaches to Challenge #3. We do not address Challenge #1; rather the existing reductions from general directed graphs to Eulerian ones allows us to focus on Eulerian digraphs without loss of generality.

Specifically, we introduce a novel definition of approximation for asymmetric matrices, which we call *singular-value approximation* (or *SV approximation*). For simplicity in the introduction, we focus on the case of regular directed graphs, i.e. directed graphs where for some value $d \geq 0$, every vertex has in-degree d and out-degree d . (In the case of digraphs with nonnegative edge weights, we obtain the in- and out-degrees by summing the in-coming or out-going edge weights at each vertex.) However, all of our results generalize to Eulerian digraphs and some generalize to wider classes of complex matrices. To introduce SV approximation, let \mathbf{A} be the adjacency matrix of a d -regular digraph, i.e. \mathbf{A} is a nonnegative real matrix where every row and column sum equals d . Then the (in- and out-) degree matrix is simply $d\mathbf{I}$. Dividing by d , it is equivalent to study approximation of the *random-walk matrix* $\mathbf{W} = \mathbf{A}/d$, which is doubly stochastic, and has degree matrix \mathbf{I} .

Definition 1.1 (SV approximation for doubly stochastic matrices). For doubly stochastic matrices $\mathbf{W}, \widetilde{\mathbf{W}} \in \mathbb{R}^{n \times n}$ we say that $\widetilde{\mathbf{W}}$ is an ε -*singular-value (SV) approximation* of \mathbf{W} , written $\widetilde{\mathbf{W}} \approx_{\varepsilon}^{sv} \mathbf{W}$, if for all vectors $x, y \in \mathbb{R}^n$, we have

$$\left| x^{\top} (\widetilde{\mathbf{W}} - \mathbf{W}) y \right| \leq \frac{\varepsilon}{4} \cdot \left[x^{\top} (\mathbf{I} - \mathbf{W}\mathbf{W}^{\top}) x + y^{\top} (\mathbf{I} - \mathbf{W}^{\top} \mathbf{W}) y \right]. \quad (1)$$

This formulation of SV-approximation is one of several equivalent formulations we provide later in the paper (Lemma 3.7). We can equivalently define SV approximation between doubly stochastic matrices by requiring Equation (1) to hold for all complex test vectors $x, y \in \mathbb{C}^n$. SV-approximation can also be defined equivalently by replacing condition (1) with

$$\left| x^{\top} (\widetilde{\mathbf{W}} - \mathbf{W}) y \right| \leq \frac{\varepsilon}{4} \cdot \sqrt{[x^{\top} (\mathbf{I} - \mathbf{W}\mathbf{W}^{\top}) x] \cdot [y^{\top} (\mathbf{I} - \mathbf{W}^{\top} \mathbf{W}) y]}. \quad (2)$$

These two formulations, (1) and (2), differ only in using the geometric mean or the arithmetic mean of the terms involving x and y on the right-hand side. The formulation in terms of the geometric mean implies the one in terms of the arithmetic mean (since the geometric mean is no larger than the arithmetic mean); the converse follows by optimizing over scalar multiples of x and y (as was done in e.g. [CKP⁺17, AKM⁺20]). Both formulations can also be rewritten more simply by noting that

$$x^{\top} (\mathbf{I} - \mathbf{W}\mathbf{W}^{\top}) x = \|x\|^2 - \|x^{\top} \mathbf{W}\|^2 \quad \text{and} \quad y^{\top} (\mathbf{I} - \mathbf{W}^{\top} \mathbf{W}) y = \|y\|^2 - \|\mathbf{W} y\|^2,$$

but the description in terms of quadratic forms will be more convenient for comparison with previous notions of approximation. In Section 3, we provide more general definitions of SV approximation which also apply to unnormalized directed Laplacians and even to complex matrices.

We prove that SV approximation is strictly stronger than previous notions of spectral approximation considered in the literature for both asymmetric and symmetric matrices, and enjoys several useful properties not possessed by the previous notions. Most notably, there is a simple black-box reduction from SV-sparsifying Eulerian directed graphs to SV-sparsifying undirected graphs; no such reduction is known for prior notions of asymmetric spectral approximation.

Furthermore, we give efficient algorithms for working with SV approximation. These include nearly linear-time algorithms for SV-sparsifying undirected and hence also Eulerian directed graphs (Theorem 1.8), as well as random-walk polynomials associated with such graphs (Theorem 1.9). We also show that a simple repeated-squaring and sparsification algorithm for solving Laplacian systems also works for Eulerian digraphs whose random-walk matrix is normal (i.e. unitarily diagonalizable), if we use SV-sparsification at each step (Theorem 1.11). Prior Laplacian solvers for Eulerian graphs are more complex. We elaborate on these results in the next several subsections.

1.2 Comparison to Previous Notions of Approximation

Let us compare Definition 1.1 to previous definitions of approximation.

Undirected spectral approximation. Let's start with the undirected case, where $\mathbf{W} = \mathbf{W}^\top$. In this case, it can be shown that we can without loss of generality restrict to $x = y \in \mathbb{R}^n$, and $\widetilde{\mathbf{W}} \overset{sv}{\approx}_\varepsilon \mathbf{W}$ requires that for all $x \in \mathbb{R}^n$,

$$\left| x^\top (\widetilde{\mathbf{W}} - \mathbf{W})x \right| \leq \frac{\varepsilon}{2} \cdot \left[x^\top (\mathbf{I} - \mathbf{W}^2)x \right]. \quad (3)$$

In contrast, the standard definition of spectral approximation (originally introduced by Spielman and Teng [ST04]), which we denote by $\widetilde{\mathbf{W}} \approx_\varepsilon \mathbf{W}$, is equivalent to requiring that for all $x \in \mathbb{R}^n$, we have

$$\left| x^\top (\widetilde{\mathbf{W}} - \mathbf{W})x \right| \leq \varepsilon \cdot \left[x^\top (\mathbf{I} - \mathbf{W})x \right]. \quad (4)$$

To compare inequalities (3) and (4), we write $x = \sum_i c_i v_i$, where v_1, \dots, v_n is an orthonormal eigenbasis for \mathbf{W} with associated eigenvalues $\lambda_1, \dots, \lambda_n$. Since \mathbf{W} is stochastic, $|\lambda_i| \leq 1$ for all $i \in [n]$. Then the right-hand side of SV inequality (3) becomes

$$\frac{\varepsilon}{2} \cdot \sum_{i \in [n]} c_i^2 \cdot (1 - \lambda_i^2),$$

whereas the right-hand side of ST inequality (4) becomes

$$\varepsilon \cdot \sum_{i \in [n]} c_i^2 \cdot (1 - \lambda_i).$$

Since each $|\lambda_i| \leq 1$, the fact that SV approximation implies ST approximation then follows from

$$(1 - \lambda_i^2) = (1 - \lambda_i)(1 + \lambda_i) \leq 2(1 - \lambda_i).$$

However, we also see that inequality (3) can be much stronger than inequality (4) when \mathbf{W} has eigenvalues λ_i close to -1 (e.g. in a bipartite graph with poor expansion) because then $1 - \lambda_i^2$ is close to 0, but $1 - \lambda_i$ is bigger than 2. More generally, inequality (3) requires that $\widetilde{\mathbf{W}}$ approximates \mathbf{W} very well on every test vector x that is concentrated on the eigenvectors whose eigenvalues have *magnitude* close to 1, whereas inequality (4) only requires close approximation on the (signed) eigenvalues that are close to 1.

We remark that another way of ensuring $\widetilde{\mathbf{W}}$ preserves unit singular values is to replace \mathbf{W}^2 in the SV inequality (3) with the matrix $|\mathbf{W}|$ where we replace all eigenvalues of \mathbf{W} with their absolute value rather than their square,¹ so that we have:

$$x^\top (\mathbf{I} - |\mathbf{W}|)x = \sum_{i \in [n]} c_i^2 \cdot (1 - |\lambda_i|).$$

Using $|\mathbf{W}|$ instead of \mathbf{W}^2 results in an equivalent definition up to a factor of 2 in ε , and \mathbf{W}^2 turns out to be convenient to work with.² This viewpoint also explains why we stop at \mathbf{W}^2 in the definition and don't explicitly use higher powers; it is simply a convenient proxy for $|\mathbf{W}|$, which captures all powers. Indeed, for all $k \in \mathbb{N}$

$$\mathbf{I} - \mathbf{W}^2 \preceq 2 \cdot (1 - |\mathbf{W}|) \preceq 2 \cdot (\mathbf{I} - \mathbf{W}^k).$$

Directed spectral approximation. Turning to previous notions of spectral approximation for directed graphs, *standard approximation* [CKP⁺17] generalizes the definition of Spielman and Teng [ST04] by saying $\widetilde{\mathbf{W}} \approx_\varepsilon \mathbf{W}$ if for all $x, y \in \mathbb{R}^n$

$$\left| x^\top (\widetilde{\mathbf{W}} - \mathbf{W})y \right| \leq \frac{\varepsilon}{2} \cdot \left[x^\top (\mathbf{I} - \mathbf{W})x + y^\top (\mathbf{I} - \mathbf{W})y \right]. \quad (5)$$

Equivalently, we can require that for all $x, y \in \mathbb{R}^n$,

$$\left| x^\top (\widetilde{\mathbf{W}} - \mathbf{W})y \right| \leq \varepsilon \cdot \sqrt{x^\top (\mathbf{I} - \mathbf{W})x \cdot y^\top (\mathbf{I} - \mathbf{W})y}. \quad (6)$$

It can be shown that for undirected graphs, standard approximation is equivalent to the condition of Equation 4, so we will also refer to it as standard approximation.

The use of different left and right test vectors x and y on the left-hand side is crucial for capturing the asymmetric information in $\widetilde{\mathbf{W}}$ and \mathbf{W} . As before, if x or y is concentrated on eigenvectors of \mathbf{W} whose eigenvalues are close to 1, then the right-hand side of ST inequality (6) is close to 0 and $\widetilde{\mathbf{W}}$ must approximate \mathbf{W} very well. However, like the standard undirected ST inequality (4), not much is required on eigenvalues near -1. Moreover, asymmetric matrices can have eigenvalues that are not real and are equal to or close to complex numbers of magnitude 1. For example, the eigenvalues of a directed n -cycle are the complex n 'th roots of unity.

To address this issue, *unit-circle (UC) approximation* [AKM⁺20], written $\widetilde{\mathbf{W}} \overset{\circ}{\approx}_\varepsilon \mathbf{W}$, requires that for all *complex* test vectors $x, y \in \mathbb{C}^n$, we have

$$\left| x^* (\widetilde{\mathbf{W}} - \mathbf{W})y \right| \leq \frac{\varepsilon}{2} \cdot \left[\|x\|^2 + \|y\|^2 - |x^* \mathbf{W}x + y^* \mathbf{W}y| \right]. \quad (7)$$

That is, we take the complex magnitude of the terms involving \mathbf{W} on the right-hand side of ST inequality (5). That way, if x and y are concentrated on eigenvectors of \mathbf{W} that have eigenvalue near some complex number μ of magnitude 1, we require that $\widetilde{\mathbf{W}}$ approximates \mathbf{W} very well. For example, consider the case where \mathbf{W} is normal, i.e. has an orthonormal basis of complex eigenvectors v_1, \dots, v_n and with corresponding complex eigenvalues $\lambda_1, \dots, \lambda_n$. Then if we write $x = \sum_i c_i v_i$ and $y = \sum_i d_i v_i$, the right-hand side of UC inequality (7) becomes:

$$\frac{\varepsilon}{2} \cdot \sum_{i \in [n]} (|c_i|^2 + |d_i|^2) - \left| \sum_{i \in [n]} (|c_i|^2 + |d_i|^2) \cdot \lambda_i \right|. \quad (8)$$

¹Another way of describing $|\mathbf{W}|$ is as the psd square root of the psd matrix \mathbf{W}^2 .

² $|\mathbf{W}| = (\mathbf{W}^2)^{1/2}$, i.e. $|\mathbf{W}|$ is the PSD square root of \mathbf{W}^2 . In Definition 1.1, we could similarly replace $\mathbf{W}\mathbf{W}^\top$ and $\mathbf{W}^\top \mathbf{W}$ with their PSD square roots and obtain a definition that is equivalent up to a factor of 2 in ε .

If x and y are concentrated on eigenvalues $\lambda_i \approx \mu$ where $|\mu| = 1$, then this expression will be close to 0. Unit-circle approximation has valuable properties not enjoyed by standard approximation, in particular being preserved under powering: If $\widetilde{\mathbf{W}}$ is an ε -UC approximation of \mathbf{W} , then for every positive integer k , $\widetilde{\mathbf{W}}^k$ is an $O(\varepsilon)$ -UC approximation of \mathbf{W}^k ; note that the quality of approximation does not degrade with k . This property was crucial for the results of [AKM⁺20].

However, UC approximation has two limitations compared to SV approximation. First, UC expression (8) is only small if x and/or y is concentrated on eigenvalues that are all close to the *same* point μ on the complex unit circle. Even in the undirected case, if x and y are mixtures of eigenvectors with eigenvalue close to 1 and eigenvalue close to -1, then there will be cancellations in the second term of UC expression (8) and the result will not be small. Second, the spectral properties of asymmetric matrices are better captured by *singular values* than eigenvalues, since singular values treat the domain and codomain as distinct.

To see how SV approximation addresses these limitations, let $\sigma_1, \dots, \sigma_n \geq 0$ be the singular values of \mathbf{W} , let $u_1, u_2, \dots, u_n \in \mathbb{C}^n$ the corresponding left-singular vectors of \mathbf{W} , and let $v_1, \dots, v_n \in \mathbb{C}^n$ the corresponding right-singular vectors. If we write $x = \sum_i c_i u_i$ and $y = \sum_i d_i v_i$, then the right-hand side of SV inequality (1) becomes:

$$\frac{\varepsilon}{4} \cdot \left[\sum_{i \in [n]} (|c_i|^2 + |d_i|^2) \cdot (1 - \sigma_i^2) \right]. \quad (9)$$

Consequently, SV-approximation requires high-quality approximation if x is concentrated on left-singular vectors of singular value close to 1 and/or y is concentrated on right-singular vectors of singular value close to 1. (For the “or” interpretation, use the formulation of SV approximation in terms of inequality (2).) To compare with UC expression (8), let us consider what happens with a *normal* matrix, where $u_i = v_i$ and $\sigma_i = |\lambda_i|$. In this case, SV expression (9) amounts to bringing the absolute value of UC expression (8) inside the summation (and squaring, which only makes a factor of 2 difference), to avoid cancellations between eigenvalues of different phases.

Furthermore, for non-normal matrices, SV approximation retains the asymmetry of \mathbf{W} even on the right-hand side, by always using x on the left of \mathbf{W} (thus relating to its decomposition into left singular vectors) and y on the right of \mathbf{W} (thus relating to its decomposition into right singular vectors). Indeed, this feature allows us to even extend the definition of SV approximation to non-square matrices. (See Section 3.)

Following the above intuitions, we prove that SV approximation is indeed strictly stronger than the previous notions of approximation:

Theorem 1.2. *For all doubly stochastic \mathbf{W} and $\widetilde{\mathbf{W}}$, if $\widetilde{\mathbf{W}} \overset{sv}{\approx}_\varepsilon \mathbf{W}$, then $\widetilde{\mathbf{W}} \overset{\circ}{\approx}_\varepsilon \mathbf{W}$ (and hence $\widetilde{\mathbf{W}} \approx_\varepsilon \mathbf{W}$). On the other hand, for every $\varepsilon \in (0, 1)$ and $\varepsilon' \in (\varepsilon, \infty)$, there exist symmetric $\mathbf{W}, \widetilde{\mathbf{W}} \in \mathbb{R}^{2 \times 2}$ of spectral norm at most 1 such that $\widetilde{\mathbf{W}} \overset{\circ}{\approx}_\varepsilon \mathbf{W}$ but it is not the case that $\widetilde{\mathbf{W}} \overset{svn}{\approx}_{\varepsilon'} \mathbf{W}$.*

The separation above relies on a generalization of the definition of SV approximation to matrices that can have negative entries, given in Section 3. It would be interesting to extend the separation to matrices corresponding to graphs, and even doubly stochastic matrices. We note that in [AKM⁺20] UC approximation was separated from standard approximation even for doubly stochastic matrices, so SV approximation is also strictly stronger than standard approximation, even for doubly stochastic matrices.

Other Work on SV Approximation. The definition of SV approximation and some of our results on it (obtained in collaboration between Jack Murtagh and the authors) were first presented in

the first author’s PhD thesis [Ahm20] in August 2020. Independently, Kelley [Kel21] used a variant of SV approximation to present an alternative proof of a result of [HPV21], who used unit-circle approximation to prove that the Impagliazzo–Nisan–Wigderson pseudorandom generator [INW94] fools permutation branching programs. Golowich and Vadhan [GV22] used SV approximation and some of our results (presented in the aforementioned thesis) to prove new pseudorandomness properties of expander walks against permutation branching programs. Most recently, Chen, Lyu, Tal, and Wu [CLTW22] have used a form of SV approximation to present alternative proofs of the results of [AKM⁺20, PV21].

1.3 Properties of SV Approximation

SV approximation enjoys a number of novel properties not known to be possessed by previous notions of spectral approximation. Most striking is the fact that directed approximation reduces to undirected approximation. To formulate this, we define the symmetric lift of a matrix:

Definition 1.3. Given $\mathbf{W} \in \mathbb{C}^{m \times n}$, let the *symmetric lift* of \mathbf{W} be defined as

$$\text{sift}(\mathbf{W}) \stackrel{\text{def}}{=} \begin{bmatrix} \mathbf{0}^{n \times n} & \mathbf{W}^* \\ \mathbf{W} & \mathbf{0}^{m \times m} \end{bmatrix}.$$

Graph theoretically, the symmetric lift of \mathbf{W} is the following standard operation: Given our directed graph G on n vertices with random-walk matrix \mathbf{W} , we lift G to an undirected bipartite graph H with n vertices on each side, where we connect left-vertex i to right-vertex j if there is a directed edge from i to j in G . Then $\text{sift}(\mathbf{W})$ is the random-walk matrix of H .

Theorem 1.4. *Let \mathbf{W} and $\widetilde{\mathbf{W}}$ be doubly stochastic matrices. Then $\mathbf{W} \stackrel{sv}{\approx}_\varepsilon \widetilde{\mathbf{W}}$ if and only if $\text{sift}(\widetilde{\mathbf{W}}) \stackrel{sv}{\approx}_\varepsilon \text{sift}(\mathbf{W})$.*

Thus, for the first time (as far as we know), sparsification of directed graphs reduces directly to sparsification of undirected graphs. It would be very interesting to obtain a similar reduction for other algorithmic problems in spectral graph theory, such as solving Laplacian systems.

Another novel property of SV approximation is that it is preserved under products of random-walk matrices:

Theorem 1.5. *Let $\mathbf{W}_1, \dots, \mathbf{W}_k$ and $\widetilde{\mathbf{W}}_1, \dots, \widetilde{\mathbf{W}}_k$ be doubly stochastic matrices such that $\widetilde{\mathbf{W}}_i \stackrel{sv}{\approx}_\varepsilon \mathbf{W}_i$ for each $i \in [k]$. Then $\widetilde{\mathbf{W}}_1 \widetilde{\mathbf{W}}_2 \cdots \widetilde{\mathbf{W}}_k \stackrel{sv}{\approx}_{\varepsilon + O(\varepsilon^2)} \mathbf{W}_1 \mathbf{W}_2 \cdots \mathbf{W}_k$.*

Notably the approximation error does not grow with the number k of matrices being multiplied. The analogous property was known for UC approximation only in the special case of powering, i.e. $\mathbf{W}_1 = \mathbf{W}_2 = \cdots = \mathbf{W}_k$ and $\widetilde{\mathbf{W}}_1 = \widetilde{\mathbf{W}}_2 = \cdots = \widetilde{\mathbf{W}}_k$.

In addition, SV approximation is preserved under multiplication on the left and right by arbitrary matrices of bounded spectral norm. Indeed, it can be seen as the “closure” of standard approximation under this operation (up to a factor of 2).

Theorem 1.6.

1. *Let \mathbf{W} and $\widetilde{\mathbf{W}}$ be doubly stochastic matrices such that $\widetilde{\mathbf{W}} \stackrel{sv}{\approx}_\varepsilon \mathbf{W}$. Then for all complex matrices \mathbf{U} and \mathbf{V} of spectral norm at most 1, we have $\mathbf{U} \widetilde{\mathbf{W}} \mathbf{V} \stackrel{sv}{\approx}_\varepsilon \mathbf{U} \mathbf{W} \mathbf{V}$, and hence $\mathbf{U} \widetilde{\mathbf{W}} \mathbf{V} \approx_\varepsilon \mathbf{U} \mathbf{W} \mathbf{V}$.*
2. *Let \mathbf{W} and $\widetilde{\mathbf{W}}$ be doubly stochastic matrices such that for all complex matrices \mathbf{U} and \mathbf{V} of spectral norm at most 1, we have $\mathbf{U} \widetilde{\mathbf{W}} \mathbf{V} \approx_\varepsilon \mathbf{U} \mathbf{W} \mathbf{V}$. Then $\widetilde{\mathbf{W}} \stackrel{sv}{\approx}_{2\varepsilon} \mathbf{W}$.*

Since $\mathbf{U}\mathbf{W}\mathbf{V}$ and $\mathbf{U}\widetilde{\mathbf{W}}\mathbf{V}$ need not be doubly stochastic matrices, Theorem 1.6 uses the generalization of SV approximation to more general matrices, which can be found in Section 3.

Recall that standard spectral sparsifiers [ST04] are also *cut sparsifiers* [BK00]. That is, if \widetilde{G} is an ε -approximation of G , then for every set S of vertices, the weight of the cut S in \widetilde{G} is within a $(1 \pm \varepsilon)$ factor of the weight of S in G . Indeed, if we take the test vector x to be the characteristic vector of the set S in inequality (4), we obtain

$$\left| \Pr_{(i,j) \sim \mu_{\text{edge}}} [i \in S, j \notin S] - \Pr_{(i,j) \sim \widetilde{\mu}_{\text{edge}}} [i \in S, j \notin S] \right| \leq \varepsilon \cdot \Pr_{(i,j) \sim \mu_{\text{edge}}} [i \in S, j \notin S], \quad (10)$$

where $\mu_{\text{edge}} \stackrel{\text{def}}{=} \mu_{\text{edge}}(\mathbf{W})$ (resp. $\widetilde{\mu}_{\text{edge}} \stackrel{\text{def}}{=} \widetilde{\mu}_{\text{edge}}(\widetilde{\mathbf{W}})$) denotes a random edge according to the stationary distribution of \mathbf{W} (resp. $\widetilde{\mathbf{W}}$), namely pick $i \in [n]$ uniformly at random and obtain j as a random neighbor of i using \mathbf{W} (resp. $\widetilde{\mathbf{W}}$). Note that $\Pr_{(i,j) \sim \mu_{\text{edge}}} [i \in S, j \notin S]$ is simply a normalized version of the weight of the cut (S, S^c) ; namely it equals the sum of the edge weights leaving S divided the sum of all edge weights in the graph.

Similarly, we can obtain a combinatorial consequence of SV approximation, by taking x to be a characteristic vector of a set S of vertices and taking y to be a characteristic vector of a set T of vertices. This yields:

Proposition 1.7. *Let $\widetilde{\mathbf{W}}$ and \mathbf{W} be doubly stochastic $n \times n$ random-walk matrices and suppose that $\widetilde{\mathbf{W}} \stackrel{sv}{\approx}_{\varepsilon} \mathbf{W}$. Then for every two subsets $S, T \subseteq [n]$, we have*

$$\left| \Pr_{(i,j) \sim \mu_{\text{edge}}} [i \in S, j \in T] - \Pr_{(i,j) \sim \widetilde{\mu}_{\text{edge}}} [i \in S, j \in T] \right| \leq \frac{\varepsilon}{2} \cdot \sqrt{\Pr_{(i,j,k) \sim \mu_{\text{fb}}} [i \in S, k \notin S] \cdot \Pr_{(i,j,k) \sim \mu_{\text{fb}}} [i \in T, k \notin T]},$$

where:

- $\mu_{\text{edge}} \stackrel{\text{def}}{=} \mu_{\text{edge}}(\mathbf{W})$ (resp. $\widetilde{\mu}_{\text{edge}} \stackrel{\text{def}}{=} \widetilde{\mu}_{\text{edge}}(\widetilde{\mathbf{W}})$) denotes a random edge according to the stationary distribution of \mathbf{W} (resp. $\widetilde{\mathbf{W}}$), namely pick $i \in [n]$ uniformly at random and obtain j as a random neighbor of i using \mathbf{W} (resp. $\widetilde{\mathbf{W}}$), and
- $\mu_{\text{fb}} \stackrel{\text{def}}{=} \mu_{\text{fb}}(\mathbf{W})$ denotes a random forward-backward walk, namely pick $i \in [n]$ uniformly at random, obtain j as a random neighbor of i using \mathbf{W} , and k as a random neighbor of j using \mathbf{W}^{\top} .

Let us interpret Proposition 1.7. First, consider the case that $\mathbf{W} = \mathbf{J}$, the matrix with every entry equal to $1/n$ (the random-walk matrix for the complete graph with self-loops). Then $(i, j) \sim \mu_{\text{edge}}$ and $(i, j, k) \sim \mu_{\text{fb}}$ are all uniform and independent vertices, so Proposition 1.7 says:

$$\left| \mu(S) \cdot \mu(T) - \Pr_{(i,j) \sim \mu_{\text{edge}}} [i \in S, j \in T] \right| \leq \frac{\varepsilon}{2} \cdot \sqrt{\mu(S) \cdot (1 - \mu(S)) \cdot \mu(T) \cdot (1 - \mu(T))},$$

where $\mu(S) = |S|/n$ and $\mu(T) = |T|/n$ are the stationary probabilities of S and T , respectively. This amounts to a restatement of the Expander Mixing Lemma ([Vad12, Lemma 4.15]); indeed $\widetilde{\mathbf{W}} \stackrel{sv}{\approx}_{\varepsilon} \mathbf{J}$ if and only if $\widetilde{\mathbf{W}}$ is a spectral expander with all nontrivial singular values at most $\varepsilon/2$.

Next, let's consider the case that $T = S^c$. Then,

$$\Pr_{(i,j,k) \sim \mu_{\text{fb}}} [i \in S, k \notin S] = \Pr_{(i,j,k) \sim \mu_{\text{fb}}} [i \in T, k \notin T].$$

(In an Eulerian graph, the weight of edges crossing any cut in one direction must equal the weight of edges crossing it in the other direction.) Thus, SV approximation implies that:

$$\left| \Pr_{(i,j) \sim \mu_{\text{edge}}} [i \in S, j \notin S] - \Pr_{(i,j) \sim \tilde{\mu}_{\text{edge}}} [i \in S, j \notin S] \right| \leq \frac{\varepsilon}{2} \cdot \Pr_{(i,j,k) \sim \mu_{\text{fb}}} [i \in S, k \notin S]. \quad (11)$$

We claim that (11) is stronger than the standard notion of a cut approximator (10). Indeed,

$$\Pr_{(i,j,k) \sim \mu_{\text{fb}}} [i \in S, k \notin S] \leq \Pr_{(i,j,k) \sim \mu_{\text{fb}}} [i \in S, j \notin S] + \Pr_{(i,j,k) \sim \mu_{\text{fb}}} [j \in S, k \notin S] = 2 \Pr_{(i,j) \sim \mu_{\text{edge}}} [i \in S, j \notin S]. \quad (12)$$

Similarly, we have

$$\Pr_{(i,j,k) \sim \mu_{\text{fb}}} [i \in S, k \notin S] \leq \Pr_{(i,j) \sim \mu_{\text{edge}}} [i \in S, j \in S] + \Pr_{(i,j) \sim \mu_{\text{edge}}} [i \notin S, j \notin S] \quad (13)$$

Thus, we conclude that an SV-approximator not only approximates every cut to within a small additive error that is scaled by the weight of the cut edges (as in (10)), but also scaled by the weight of the *uncut* edges.

1.4 Algorithmic Results

Even though SV approximation is stronger than previously considered notions of spectral approximation, we show that it still admits sparsification:

Theorem 1.8. *Given a regular directed graph G with n vertices and m edges, integer edge weights in $[0, U]$, and random-walk matrix \mathbf{W} , and $\varepsilon > 0$*

1. *There exists a weighted graph \tilde{G} with at most $O(n\varepsilon^{-2} \log(nU) \log^{11} n)$ edges such that its random-walk matrix $\tilde{\mathbf{W}}$ satisfies $\tilde{\mathbf{W}} \stackrel{sv}{\approx}_{\varepsilon} \mathbf{W}$.*
2. *There is a randomized nearly-linear time algorithm that whp outputs a weighted graph \tilde{G} with at most $O(n\varepsilon^{-2} \log(nU) \log^{19} n)$ edges such that its random-walk matrix $\tilde{\mathbf{W}}$ satisfies $\tilde{\mathbf{W}} \stackrel{sv}{\approx}_{\varepsilon} \mathbf{W}$.*

A more general theorem that also applies to Eulerian digraphs is stated in the main body of the paper (Corollaries 4.12 and 4.13). Prior to this work, it was open whether or not even UC-sparsifiers with $O(n \cdot \text{poly}(\log n, \varepsilon^{-1}))$ edges existed for all unweighted regular digraphs. Instead, it was only known how to UC-sparsify powers of a random walk matrix to bound the sparsity increase.³

By Theorem 1.4, it suffices to prove Theorem 1.8 for undirected bipartite graphs. We obtain the latter via an undirected sparsification algorithms based on expander partitioning [ST04]. It remains an open question whether algorithms based on edge sampling can yield SV approximation or unit circle approximation, even in undirected graphs. The standard approach to spectral sparsification of undirected graphs via sampling, namely keeping each edge independently with probability proportional to its effective resistance [SS08], does not work for SV or UC approximation. For example, this method does not exactly preserve degrees, which we show is necessary for SV sparsification (Lemma 3.17).⁴

However, we remark that the work of Chu, Gao, Peng, Sawlani, and Wang [CGP⁺18] does yield something closer to sparsification via *degree preserving* sampling for standard approximation [CKP⁺17] but not unit circle approximation. They show that if one has a directed graph and

³We extend this result to SV approximation (Lemma 4.15).

⁴Unlike standard spectral approximation, degrees cannot be fixed just by adding self loops; indeed, self-loops ruin bipartiteness and periodicity, which are properties that UC and SV approximation retain (as they are captured by eigenvalues like -1 or other roots of unity).

decomposes it into short “cycles” without regard for the direction of the edges on the cycle, then one can sparsify by randomly eliminating either the clockwise or counterclockwise edges on each such cycle. We build on their procedure and use it to obtain SV sparsification (and hence, unit circle) by showing that this technique obtains SV approximation, even if the cycles are not short, as long as (a) all the cycles are within expanding subgraphs, and (b) the cycles alternate between forward and backward edges. (Note that such alternating cycles in a directed graph correspond to ordinary cycles in the undirected lift given by Theorem 1.4.)

Sparsity	Approximation	Time	Subgraph?	Citation
$O(n\varepsilon^{-2} \log^c n)$	Standard	$O(m \log^c n)$	No	[CKP ⁺ 17]
$O(n\varepsilon^{-2} \log^c n)$	Standard	$O(m^{1+o(1)})$	Yes	[CGP ⁺ 18, PY19]
$O(n\varepsilon^{-2} \log^4 n)$	Standard	$O(nm)$	Yes	[CGP ⁺ 18]
$O(n\varepsilon^{-2} \log^{12} n)$	SV	Existential	Yes	Corollary 4.13
$O(n\varepsilon^{-2} \log^{20} n)$	SV	$O(m \log^7 n)$	Yes	Corollary 4.12

Our results also enable us to give the first nearly linear-time algorithm for sparsifying powers of random walk matrices of graphs. More precisely, we show the following.

Theorem 1.9 (Theorem 4.17, informally stated). *There is a randomized nearly-linear time algorithm that, given a regular digraph G with random walk matrix \mathbf{W} and any positive integer $k \leq \text{poly}(n)$, outputs a nearly linear sized SV-sparsifier of \mathbf{W}^k , provided that the second largest singular value of \mathbf{W} is at most $1 - 1/\text{poly}(n)$.*

This theorem is stated more generally for Eulerian digraphs and proven as Theorem 4.17.

A regular (or even Eulerian) digraph whose weak mixing time (i.e. the mixing time of the 1/2-lazy random walk) is at most $\text{poly}(n)$ can be made to have second largest singular value at most $1 - 1/\text{poly}(n)$ by making it $1/\text{poly}(n)$ -lazy (Lemma B.5). Thus we obtain the following corollary that allows us to unconditionally sparsify lazy random walks with even very small amounts of laziness.

Corollary 1.10 (Corollary 4.19, informally stated). *Let G be a regular digraph with $\text{poly}(n)$ weak mixing time. For any positive integer $k \leq \text{poly}(n)$ and $\gamma \geq 1/\text{poly}(n)$, we can, in nearly linear time, output a nearly linear sized SV-sparsifier of $((1 - \gamma)\mathbf{W} + \gamma\mathbf{I})^k$.*

Importantly we can take $\gamma \ll 1/k$, so a γ -lazy walk of length k is very close (e.g. in total variation distance) to an ordinary random walk of length k .

Our algorithm is relatively simple: we show that we can SV-sparsify the square of random walk matrices of regular graphs, which gives all powers of two, then multiply different such powers together to get the precise exponent we want. For example, to get a sparsifier for \mathbf{W}^7 , we multiply sparsifiers for \mathbf{W}^4 , \mathbf{W}^2 , and \mathbf{W}^1 . The use of SV approximation plays an important role in the analysis of this algorithm, because it has the property that the product of the approximations of the powers still approximates the product of the true powers (Theorem 1.5).

Finally for Challenge #3, we make progress towards simplifying the recursion and analysis of solving directed Laplacian linear system in the following way. We show that a simpler recursion, a variant of the Peng-Spielman repeated squaring based recursion (Equation (15)) and analysis suffice if the directed Laplacian is normal (i.e. unitarily diagonalizable). A solver of this form was given in the undirected case by Peng and Spielman [PS14], but the existing solvers for directed graphs are all significantly more involved. Note that this result on the squaring recursion is the only result in this paper that relies on a normality assumption, e.g. aforementioned sparsification results hold for all Eulerian directed graphs.

Theorem 1.11. For a doubly stochastic normal matrix $\mathbf{W} \in \mathbb{R}^{n \times n}$ with $\|\mathbf{W}\| \leq 1$, let $\mathbf{W} = \mathbf{W}_0, \dots, \mathbf{W}_{k-1}$ be a sequence of matrices such that for $\varepsilon \leq 1/4k$ we have

$$\mathbf{W}_i \stackrel{sv}{\approx}_\varepsilon \mathbf{W}_{i-1}^2 \quad \forall 0 < i < k, \quad (14)$$

and

$$\mathbf{P}_i = \frac{1}{2} [\mathbf{I} + (\mathbf{I} + \mathbf{W}_i)\mathbf{P}_{i+1}(\mathbf{I} + \mathbf{W}_i)] \quad \forall 0 \leq i < k \quad (15)$$

defining the Peng-Spielman squaring recursion.

Then, for a matrix \mathbf{P}_k , such that $\left\| (\mathbf{I} - \mathbf{W}^{2^k})^{\frac{1}{2}} [\mathbf{P}_k - (\mathbf{I} - \mathbf{W}^{2^k})^+] (\mathbf{I} - \mathbf{W}^{2^k})^{\frac{1}{2}} \right\| \leq O(k\varepsilon)$, we have

$$\|\mathbf{P}_0(\mathbf{I} - \mathbf{W}) - \mathbf{I}\|_{\mathbf{B}} = \left\| (\mathbf{I} - \mathbf{W})^{\frac{1}{2}} [\mathbf{P}_0 - (\mathbf{I} - \mathbf{W})^+] (\mathbf{I} - \mathbf{W})^{\frac{1}{2}} \right\| \leq O(k^2\varepsilon) \quad (16)$$

where $\mathbf{B} = ((\mathbf{I} - \mathbf{W})^{1/2})^*(\mathbf{I} - \mathbf{W})^{1/2}$.

Theorem 1.11 says that that we can compute a good *preconditioner* \mathbf{P}_0 for the Laplacian $\mathbf{I} - \mathbf{W}$ (eq. 16) by repeatedly computing SV-approximate squares (eq. 14) and use the simple recurrence (eq. 15, starting with a preconditioner \mathbf{P}_k for a sufficiently large power of \mathbf{W}). Generally, \mathbf{P}_k is easy to obtain for a large enough $k = O(\log n)$ since \mathbf{W}^{2^k} is well-approximated by a complete graph (assuming the original graph is connected and aperiodic).

1.5 Open Problems

One open problem is to determine whether or not it is possible to obtain *linear-sized* sparsifiers. Recall that undirected graphs have sparsifiers with respect to standard spectral approximation that have only $O(n/\varepsilon^2)$ nonzero edge weights [BSS12]. If this result could be extended to obtain linear-sized SV-sparsifiers of undirected graphs, we would also have linear-sized sparsifiers for directed graphs by Theorem 1.4, which would be a new result even for standard approximation [CKP⁺17].

1.6 Roadmap

The rest of the paper is organized as follows:

- Section 2 gives definitions we use throughout the paper.
- Section 3 defines SV-approximation and provides several equivalent characterizations of it which are useful when analyzing algorithms and also interesting in their own right. These results immediately imply a reduction from the directed to undirected case for SV approximation.
- Section 4 gives a direct SV sparsification algorithm based on sparsifying alternating cycles within expanders, and shows that the derandomized square gives SV square sparsifiers, and shows how to sparsify powers of random walk matrices with respect to SV approximation.
- Section 5 gives our simpler algorithm and analysis for solving Laplacian linear systems when the matrix is unitarily diagonalizable (normal).
- In Appendix A, we prove equivalences, properties, and separations, including several deferred proofs from the body of the paper.

2 Preliminaries

We begin by providing notation and definitions that we will use throughout the paper.

Complex Numbers. Let \mathbb{C} denote the set of complex numbers. For $z \stackrel{\text{def}}{=} a+bi \in \mathbb{C}$, let $z^* \stackrel{\text{def}}{=} a-bi$ and $\Re[z] \stackrel{\text{def}}{=} a$ and $\Im[z] \stackrel{\text{def}}{=} b$. Define the *magnitude* of z as $|z| \stackrel{\text{def}}{=} \sqrt{z^*z}$.

Matrix Notation. For $x \in \mathbb{C}^n$ we let $\mathbf{diag}(x) \in \mathbb{C}^{n \times n}$ denote the diagonal matrix with $[\mathbf{diag}(x)]_{i,i} = x_i$ for all $i \in [n]$. For complex matrix \mathbf{M} that is not necessarily square, we write \mathbf{M}^* to denote the conjugate transpose of \mathbf{M} and \mathbf{M}^+ to denote the Moore-Penrose pseudoinverse of \mathbf{M} . We let $\text{lker}(\mathbf{M})$ and $\text{rker}(\mathbf{M})$ be the left and right kernels of \mathbf{M} respectively. We say $\mathbf{M} \in \mathbb{C}^{n \times n}$ is *Hermitian* if $\mathbf{M}^* = \mathbf{M}$. \mathbf{M} is *positive semidefinite (PSD)* if it is Hermitian and for every vector $x \in \mathbb{C}^n$, we have $x^*\mathbf{M}x \geq 0$. \mathbf{M} is *normal* if $\mathbf{M}^*\mathbf{M} = \mathbf{M}\mathbf{M}^*$. For a square matrix \mathbf{M} , let $\mathbf{S}_\mathbf{M} = (\mathbf{M} + \mathbf{M}^*)/2$ denote its symmeterization. Note that $\mathbf{S}_\mathbf{M}$ is Hermitian, and if \mathbf{M} is a scalar then $\mathbf{S}_\mathbf{M} = \Re[\mathbf{M}]$. Given any $\mathbf{M} \in \mathbb{C}^{n \times n}$ and subsets $S, T \subset [n]$, we let $\mathbf{M}_{S,T} \in \mathbb{C}^{S \times T}$ be the submatrix of \mathbf{M} with rows specified by S and columns specified by T . We let $S^c \stackrel{\text{def}}{=} [n] \setminus S$ be the complement of S .

Schur Complements. Given an $n \times n$ matrix \mathbf{M} and $S \subset [n]$, we define the *Schur Complement of \mathbf{M} onto S* as

$$\text{SC}_S(\mathbf{M}) = \mathbf{M}_{S^c, S^c} - \mathbf{M}_{S^c, S} \mathbf{M}_{S, S}^+ \mathbf{M}_{S, S^c}.$$

Eigenvalues and Singular Values. For a symmetric matrix \mathbf{A} with nonnegative entries, let $\lambda_i(\mathbf{A})$ denote the i th largest eigenvalue of \mathbf{A} . For a matrix \mathbf{A} , let $\sigma_i(\mathbf{A}) \stackrel{\text{def}}{=} \lambda_i(\mathbf{A}^*\mathbf{A})$. Given an undirected graph G with adjacency matrix \mathbf{A} and degree matrix $\mathbf{D} \stackrel{\text{def}}{=} \mathbf{diag}(\mathbf{A}\mathbf{1})$, let $\lambda(G) \stackrel{\text{def}}{=} \lambda_2(\mathbf{D}^{+/2}\mathbf{A}\mathbf{D}^{+/2})$.

Norms. Throughout we use $\|\cdot\|$ to denote the spectral norm, where for any $\mathbf{A} \in \mathbb{C}^{n \times m}$,

$$\|\mathbf{A}\| \stackrel{\text{def}}{=} \sup_{x \in \mathbb{C}^m \setminus \{\vec{0}\}} \frac{\|\mathbf{A}x\|}{\|x\|}.$$

Löwner Order. Given Hermitian $\mathbf{A}, \mathbf{B} \in \mathbb{C}^{n \times n}$ we write $\mathbf{A} \preceq \mathbf{B}$ if $\mathbf{B} - \mathbf{A}$ is PSD, i.e. for every $x \in \mathbb{C}^n$ we have $x^*\mathbf{A}x \leq x^*\mathbf{B}x$.

3 Singular Value Approximation

In this section we formally define the notions of approximation we work with throughout the paper (Section 3.1), provide a number of equivalent definitions of SV approximation (Section 3.2), and give key properties of SV approximation (Section 3.3). All proofs in this section are deferred to their analogous subsection in Appendix A.

3.1 Matrix Approximation

First we define a general notion of matrix approximation with respect to arbitrary PSD matrices (Definition 3.1). This definition and the equivalences are a generalization of those established in [CKP⁺17].

Intuitively, we say that a matrix $\tilde{\mathbf{A}}$ is an ϵ -approximation of \mathbf{A} with respect to error matrices \mathbf{E} and \mathbf{F} if we can bound the bilinear form of their difference, i.e. $x^*(\tilde{\mathbf{A}} - \mathbf{A})y$, by the quadratic forms of x with \mathbf{E} and y with \mathbf{F} , i.e. $x^*\mathbf{E}x$ and $y^*\mathbf{F}y$. We use the term *error matrix* here to distinguish this notion of approximation from standard approximation (Definition 3.3) which we define later.

Definition 3.1 (Matrix approximation). Let $\mathbf{A}, \tilde{\mathbf{A}} \in \mathbb{C}^{m \times n}$, and let $\mathbf{E} \in \mathbb{C}^{m \times m}, \mathbf{F} \in \mathbb{C}^{n \times n}$ be PSD matrices. For $\epsilon \geq 0$, we say that $\tilde{\mathbf{A}}$ is an ϵ -approximation of \mathbf{A} with respect to error matrices \mathbf{E} and \mathbf{F} if any of the following equivalent conditions hold:

1. $\left| x^*(\tilde{\mathbf{A}} - \mathbf{A})y \right| \leq \frac{\epsilon}{2} (x^*\mathbf{E}x + y^*\mathbf{F}y)$ for all $x \in \mathbb{C}^m, y \in \mathbb{C}^n$
2. $\left| x^*(\tilde{\mathbf{A}} - \mathbf{A})y \right| \leq \epsilon \cdot \sqrt{x^*\mathbf{E}x} \cdot \sqrt{y^*\mathbf{F}y}$ for all $x \in \mathbb{C}^m, y \in \mathbb{C}^n$
3. $\left\| \mathbf{E}^{+1/2}(\tilde{\mathbf{A}} - \mathbf{A})\mathbf{F}^{+1/2} \right\| \leq \epsilon$, $\text{lker}(\tilde{\mathbf{A}} - \mathbf{A}) \supseteq \text{ker}(\mathbf{E})$, and $\text{rker}(\tilde{\mathbf{A}} - \mathbf{A}) \supseteq \text{ker}(\mathbf{F})$.

If $\mathbf{E} = \mathbf{F}$, we say that $\tilde{\mathbf{A}}$ is an ϵ -approximation of \mathbf{A} with respect to error matrix \mathbf{E} .

In the following Lemma 3.2 we specialize and simplify the equivalences presented in Definition 3.1 to Hermitian matrices.

Lemma 3.2 (Hermitian matrix approximation). Let $\mathbf{A}, \tilde{\mathbf{A}} \in \mathbb{C}^{m \times m}$ be Hermitian matrices and let $\mathbf{E} \in \mathbb{C}^{m \times m}$ be a PSD matrix. Then the following are equivalent conditions.

1. $\tilde{\mathbf{A}}$ is an ϵ -approximation of \mathbf{A} with respect to error matrix \mathbf{E} .
2. For all $x \in \mathbb{C}^m$ we have $\left| x^*(\tilde{\mathbf{A}} - \mathbf{A})x \right| \leq \epsilon \cdot (x^*\mathbf{E}x)$.
3. $-\epsilon\mathbf{E} \preceq \tilde{\mathbf{A}} - \mathbf{A} \preceq \epsilon\mathbf{E}$.

Next, for matrices $\mathbf{A}, \tilde{\mathbf{A}}, \mathbf{D} \in \mathbb{C}^{m \times m}$ we define a natural notion of approximation between $\mathbf{D} - \mathbf{A}$ and $\mathbf{D} - \tilde{\mathbf{A}}$ which we call *standard approximation* (Definition 3.3). The choice of this name is because when \mathbf{A} and $\tilde{\mathbf{A}}$ are adjacency matrices of undirected graphs with the same degrees, standard approximation coincides with spectral approximation of the Laplacian matrices of the associated graphs. Further, when \mathbf{A} and $\tilde{\mathbf{A}}$ are adjacency matrices of directed Eulerian digraphs with the same degrees then standard approximation coincides with ϵ -approximation from [CKP⁺17]. Standard approximation generalizes these two cases in a natural way even when \mathbf{D} , which we call a *degree matrix*, is not diagonal (as it was in these two cases and as we will often choose it to be). We use the term degree matrix to emphasize when we are using standard approximation (Definition 3.3) and its variants rather than (Definition 3.1).

We now present a series of definitions and equivalences and properties. For clarity, the reader can initially think of the case that $\mathbf{D} = \mathbf{I}$ and $\mathbf{A} = \mathbf{W}$ is a doubly stochastic matrix.

Definition 3.3 (Standard approximation). Let $\mathbf{A}, \tilde{\mathbf{A}} \in \mathbb{C}^{m \times m}$ and $\mathbf{D} \in \mathbb{C}^{m \times m}$ be a PSD matrix. For $\epsilon \geq 0$, we say $\tilde{\mathbf{A}}$ is a *standard ϵ -approximation of \mathbf{A} with respect to degree matrix \mathbf{D}* if $\mathbf{E} = \mathbf{D} - \mathbf{S}_{\mathbf{A}}$ is PSD and $\tilde{\mathbf{A}}$ is an ϵ -approximation of \mathbf{A} with respect to error matrix \mathbf{E} .⁵

⁵Recall by Section 2 that $\mathbf{S}_{\mathbf{A}} \stackrel{\text{def}}{=} (\mathbf{A} + \mathbf{A}^*)/2$.

With matrix approximation and standard approximation established, we can now define unit-circle (UC) approximation and singular-value (SV) approximation. UC approximation as we present it here, is a generalization of UC approximation as it was introduced for random walk matrices in [AKM⁺20]. As discussed in the introduction, SV approximation is a new notion of approximation introduced in this paper; it has a number of natural desirable properties and facilitates our results on sparsification (Section 4) and linear system solving (Section 5).

Definition 3.4 (UC approximation). Let $\mathbf{A}, \tilde{\mathbf{A}} \in \mathbb{C}^{m \times m}$ and $\mathbf{D} \in \mathbb{C}^{m \times m}$ be PSD. For $\varepsilon \geq 0$, we say $\tilde{\mathbf{A}}$ is a *unit-circle (UC) ε -approximation of \mathbf{A} with respect to degree matrix \mathbf{D}* if for every $z \in \mathbb{C}$ with $|z| = 1$, $z\tilde{\mathbf{A}}$ is a standard ε -approximation of $z\mathbf{A}$ with respect to degree matrix \mathbf{D} . If this holds, we write $\tilde{\mathbf{A}} \overset{\circ}{\approx}_{\varepsilon} \mathbf{A}$ with respect to \mathbf{D} . We omit “respect to \mathbf{D} ” when $\mathbf{D} = \mathbf{I}$.

Note that for doubly stochastic matrices $\tilde{\mathbf{W}}, \mathbf{W}$, the above definition (with $\mathbf{D} = \mathbf{I}$) corresponds to requiring for every $z \in \mathbb{C}$ and $x, y \in \mathbb{C}^n$,

$$\left| x^*(z\tilde{\mathbf{W}} - z\mathbf{W})y \right| \leq \frac{\varepsilon}{2} (\|x\|^2 + \|y\|^2 - x^*z\mathbf{W}x - y^*z\mathbf{W}y). \quad (17)$$

Optimizing over z in (17) implies Equation (7).

Definition 3.5 (SV approximation). Let $\mathbf{A}, \tilde{\mathbf{A}} \in \mathbb{C}^{m \times n}$ and let $\mathbf{D}_{\text{in}} \in \mathbb{C}^{m \times m}$ and $\mathbf{D}_{\text{out}} \in \mathbb{C}^{n \times n}$ be PSD matrices. For $\varepsilon \geq 0$, we say $\tilde{\mathbf{A}}$ is a *ε -singular-value (SV) approximation of \mathbf{A} with respect to degree matrices \mathbf{D}_{in} and \mathbf{D}_{out}* , if

1. $\ker(\mathbf{D}_{\text{in}}) \subseteq \text{lker}(\mathbf{A})$, and $\ker(\mathbf{D}_{\text{out}}) \subseteq \text{rker}(\mathbf{A})$,
2. $\mathbf{E} = \mathbf{D}_{\text{in}} - \mathbf{A}\mathbf{D}_{\text{out}}^+\mathbf{A}^*$ and $\mathbf{F} = \mathbf{D}_{\text{out}} - \mathbf{A}^*\mathbf{D}_{\text{in}}^+\mathbf{A}$ are PSD, and
3. $\tilde{\mathbf{A}}$ is an $\varepsilon/2$ -approximation of \mathbf{A} with respect to error matrices \mathbf{E} and \mathbf{F} .

If this holds, we write $\tilde{\mathbf{A}} \overset{\mathbf{D}_{\text{in}}, \mathbf{D}_{\text{out}}}{\approx}_{\varepsilon} \mathbf{A}$. If $\mathbf{A} \in \mathbb{R}_{\geq 0}^{n \times m}$ and $\mathbf{D}_{\text{in}} = \mathbf{diag}(\mathbf{A}\vec{\mathbf{1}}_n)$, $\mathbf{D}_{\text{out}} = \mathbf{diag}(\vec{\mathbf{1}}_m^{\top}\mathbf{A})$, then we write $\tilde{\mathbf{A}} \overset{sv}{\approx}_{\varepsilon} \mathbf{A}$. If $\mathbf{D}_{\text{in}} = \mathbf{I}_m$ and $\mathbf{D}_{\text{out}} = \mathbf{I}_n$, then we write $\tilde{\mathbf{A}} \overset{svn}{\approx}_{\varepsilon} \mathbf{A}$, which we call *normalized SV approximation*. If $m = n$ and $\mathbf{D}_{\text{in}} = \mathbf{D}_{\text{out}} = \mathbf{D}$, then we write $\tilde{\mathbf{A}} \overset{\mathbf{D}}{\approx}_{\varepsilon} \mathbf{A}$.

Note that normalized SV approximation does not require the relevant matrices to be non-negative, whereas $\tilde{\mathbf{A}} \overset{sv}{\approx}_{\varepsilon} \mathbf{A}$ is only defined for real non-negative matrices (such as the adjacency matrices of graphs). The maximally general definition captures both cases, so we will prove properties with respect to this notion and note their implications for the specialized notions.

3.2 Equivalent definitions of SV approximation

Here we give several equivalent formulations of SV approximation. We first give conditions under which the error matrices \mathbf{E} and \mathbf{F} in the definition of SV approximation are PSD.

Lemma 3.6 (Conditions for SV approximation to be defined). *Let $\mathbf{A}, \tilde{\mathbf{A}} \in \mathbb{C}^{m \times n}$, and let $\mathbf{D}_{\text{in}} \in \mathbb{C}^{m \times m}$ and $\mathbf{D}_{\text{out}} \in \mathbb{C}^{n \times n}$ be PSD matrices such that $\ker(\mathbf{D}_{\text{in}}) \subseteq \text{lker}(\mathbf{A})$ and $\ker(\mathbf{D}_{\text{out}}) \subseteq \text{rker}(\mathbf{A})$. Then the following are equivalent:*

1. $\sigma_{\max}(\mathbf{D}_{\text{in}}^{+/2}\mathbf{A}\mathbf{D}_{\text{out}}^{+/2}) \leq 1$.
2. $\mathbf{D}_{\text{in}} - \mathbf{A}\mathbf{D}_{\text{out}}^+\mathbf{A}^*$ is PSD.

3. $\mathbf{D}_{\text{out}} - \mathbf{A}^* \mathbf{D}_{\text{in}}^+ \mathbf{A}$ is PSD.

4. For some scalar $z \in \mathbb{C}$ with $|z| = 1$, $\begin{bmatrix} \mathbf{D}_{\text{in}} & z\mathbf{A} \\ z^* \mathbf{A}^* & \mathbf{D}_{\text{out}} \end{bmatrix}$ is PSD.

5. For every scalar $z \in \mathbb{C}$ with $|z| \leq 1$, $\begin{bmatrix} \mathbf{D}_{\text{in}} & z\mathbf{A} \\ z^* \mathbf{A}^* & \mathbf{D}_{\text{out}} \end{bmatrix}$ is PSD.

Suppose further that $\mathbf{D}_{\text{in}} = \mathbf{diag}(d_{\text{in}})$ and $\mathbf{D}_{\text{out}} = \mathbf{diag}(d_{\text{out}})$ for $d_{\text{in}} \in \mathbb{R}_{\geq 0}^m$, $d_{\text{out}} \in \mathbb{R}_{\geq 0}^n$. Then Condition 1 below implies Condition 2 below, which implies Conditions 1–5 above.

1. \mathbf{A} is nonnegative, $d_{\text{out}} = \mathbf{A} \bar{\mathbf{1}}_n$, $d_{\text{in}} = \bar{\mathbf{1}}_m^T \mathbf{A}$.

2. For all $i \in [n]$, $(d_{\text{out}})_i \geq \sum_j |\mathbf{A}_{i,j}|$, and for all $j \in [m]$, $(d_{\text{in}})_j \geq \sum_i |\mathbf{A}_{i,j}|$.

Next we give several equivalent definitions of SV approximation.

Lemma 3.7 (Equivalent formulations of SV approximation). *Let $\mathbf{A}, \tilde{\mathbf{A}} \in \mathbb{C}^{m \times n}$ and let $\mathbf{D}_{\text{in}} \in \mathbb{C}^{m \times m}$ and $\mathbf{D}_{\text{out}} \in \mathbb{C}^{n \times n}$ be PSD matrices. Then the following are equivalent*

1. $\tilde{\mathbf{A}} \stackrel{\mathbf{D}_{\text{in}}, \mathbf{D}_{\text{out}}}{\approx_\varepsilon} \mathbf{A}$.

2. $\text{sift}(\tilde{\mathbf{A}}) \stackrel{\mathbf{D}}{\approx_\varepsilon} \text{sift}(\tilde{\mathbf{B}})$, where

$$\text{sift}(\tilde{\mathbf{A}}) = \begin{bmatrix} \mathbf{0}^{m \times m} & \tilde{\mathbf{A}} \\ \tilde{\mathbf{A}}^* & \mathbf{0}^{n \times n} \end{bmatrix}, \text{sift}(\mathbf{A}) = \begin{bmatrix} \mathbf{0}^{m \times m} & \mathbf{A} \\ \mathbf{A}^* & \mathbf{0}^{n \times n} \end{bmatrix}, \mathbf{D} = \begin{bmatrix} \mathbf{D}_{\text{in}} & \mathbf{0}^{m \times n} \\ \mathbf{0}^{n \times m} & \mathbf{D}_{\text{out}} \end{bmatrix}.$$

3. For some scalar $z \in \mathbb{C}$ with $|z| = 1$, $\tilde{\mathbf{C}}$ is $\varepsilon/2$ -approximation of \mathbf{C} with respect to error matrix \mathbf{E} , where

$$\mathbf{C} = \begin{bmatrix} \mathbf{0}^{m \times m} & z\mathbf{A} \\ \mathbf{0}^{n \times m} & \mathbf{0}^{n \times n} \end{bmatrix}, \tilde{\mathbf{C}} = \begin{bmatrix} \mathbf{0}^{m \times m} & z\tilde{\mathbf{A}} \\ \mathbf{0}^{n \times m} & \mathbf{0}^{n \times n} \end{bmatrix}, \mathbf{E} = \begin{bmatrix} \mathbf{D}_{\text{in}} & z\mathbf{A} \\ z^* \mathbf{A}^* & \mathbf{D}_{\text{out}} \end{bmatrix}$$

4. Item 3 holds for every $z \in \mathbb{C}$ such that $|z| \leq 1$.

Each formulation of SV approximation in Lemma 3.7 has useful properties. Item 2 implies that SV approximation between directed graphs is equivalent to a natural related statement between *undirected* graphs, which we use for sparsification (See Section 4). A version of Item 2 is not known to hold for prior definitions of approximation between directed graphs, such as standard approximation and unit circle approximation. Item 3 and Item 4 have an error matrix \mathbf{E} that is linear in \mathbf{A} , \mathbf{D}_{in} , and \mathbf{D}_{out} , which enables short proofs of properties such as summability. In addition, Item 3 and Item 4 characterize SV approximation of \mathbf{A} in terms of ε -approximation of the ‘‘asymmetric lift’’ of \mathbf{A} with respect to its symmetrization. This enables us to leverage results developed for ε -approximation, such as preservation under Schur complements (Theorem A.2).

We next note the relation between SV and normalized SV approximation.

Lemma 3.8. *Let $\mathbf{A}, \tilde{\mathbf{A}} \in \mathbb{C}^{m \times n}$, and let $\mathbf{D}_{\text{in}} \in \mathbb{C}^{m \times m}$ and $\mathbf{D}_{\text{out}} \in \mathbb{C}^{n \times n}$ be PSD matrices such that $\ker(\mathbf{D}_{\text{in}}) \subseteq \text{lker}(\mathbf{A})$, and $\ker(\mathbf{D}_{\text{out}}) \subseteq \text{rker}(\mathbf{A})$. Let $\mathbf{N} = \mathbf{D}_{\text{in}}^{+/2} \mathbf{A} \mathbf{D}_{\text{out}}^{+/2}$ and $\tilde{\mathbf{N}} = \mathbf{D}_{\text{in}}^{+/2} \tilde{\mathbf{A}} \mathbf{D}_{\text{out}}^{+/2}$. Then $\tilde{\mathbf{A}} \stackrel{\mathbf{D}_{\text{in}}, \mathbf{D}_{\text{out}}}{\approx_\varepsilon} \mathbf{A}$ if and only if $\tilde{\mathbf{N}} \stackrel{sv}{\approx_\varepsilon} \mathbf{N}$.*

Note that if \mathbf{A} is nonnegative and $\mathbf{D}_{\text{in}} = \text{diag}(\mathbf{A}\vec{1}_n)$, $\mathbf{D}_{\text{out}} = \text{diag}(\vec{1}_m^\top \mathbf{A})$, then \mathbf{N} and $\tilde{\mathbf{N}}$ are the normalized adjacency matrices of \mathbf{A} and $\tilde{\mathbf{A}}$. Furthermore, when $\mathbf{A}, \tilde{\mathbf{A}}$ are the adjacency matrices of regular digraphs, then \mathbf{N} and $\tilde{\mathbf{N}}$ are the random-walk matrices of \mathbf{A} and $\tilde{\mathbf{A}}$ respectively. If instead \mathbf{A} and $\tilde{\mathbf{A}}$ are the adjacency matrices of Eulerian digraphs, we obtain that \mathbf{N} and $\tilde{\mathbf{N}}$ are similar to the random-walk matrices of \mathbf{A} and $\tilde{\mathbf{A}}$.

Normalized SV approximation is implied by standard $\epsilon/2$ -approximation with respect to the original degree matrix holding for all unitary multiples of the adjacency matrix.

Theorem 3.9 (Unitary transformation characterization of SV-approximation). *For $\tilde{\mathbf{N}}, \mathbf{N} \in \mathbb{C}^{n \times n}$, we have that $\tilde{\mathbf{N}} \overset{svn}{\approx}_\epsilon \mathbf{N}$ if for every pair of unitary matrices \mathbf{U}, \mathbf{V} , $\mathbf{U}\tilde{\mathbf{N}}\mathbf{V}$ is a standard $\epsilon/2$ -approximation of $\mathbf{U}\mathbf{N}\mathbf{V}$ with respect to degree matrix \mathbf{I} . Moreover, if $\tilde{\mathbf{N}} \overset{svn}{\approx}_\epsilon \mathbf{N}$ then for every pair of matrices \mathbf{U}, \mathbf{V} satisfying $\|\mathbf{U}\| \leq 1, \|\mathbf{V}\| \leq 1$, we have that $\mathbf{U}\tilde{\mathbf{N}}\mathbf{V} \overset{svn}{\approx}_\epsilon \mathbf{U}\mathbf{N}\mathbf{V}$, and hence $\mathbf{U}\tilde{\mathbf{N}}\mathbf{V}$ is a standard ϵ -approximation of $\mathbf{U}\mathbf{N}\mathbf{V}$ with respect to degree matrix \mathbf{I} .*

An important corollary is that SV approximation between adjacency matrices is preserved under multiplication on each side of the adjacency matrix by (possibly different) permutations:

Corollary 3.10 (SV preservation under multiplication by permutation matrices). *Let $\mathbf{A}, \tilde{\mathbf{A}} \in \mathbb{C}_{>0}^{m \times n}$ and suppose $\tilde{\mathbf{A}} \overset{\mathbf{D}_{\text{in}}, \mathbf{D}_{\text{out}}}{\approx}_\epsilon \mathbf{A}$. Let \mathbf{U}, \mathbf{V} be arbitrary permutation matrices. Then $\mathbf{U}\tilde{\mathbf{A}}\mathbf{V} \overset{\mathbf{D}'_{\text{in}}, \mathbf{D}'_{\text{out}}}{\approx}_\epsilon \mathbf{U}\mathbf{A}\mathbf{V}$ where $\mathbf{D}'_{\text{in}} = \mathbf{U}\mathbf{D}_{\text{in}}\mathbf{U}^*$ and $\mathbf{D}'_{\text{out}} = \mathbf{V}^*\mathbf{D}_{\text{out}}\mathbf{V}$. Consequently, if $\tilde{\mathbf{A}} \overset{sv}{\approx}_\epsilon \mathbf{A}$, then $\mathbf{U}\tilde{\mathbf{A}}\mathbf{V} \overset{sv}{\approx}_\epsilon \mathbf{U}\mathbf{A}\mathbf{V}$ and if $\tilde{\mathbf{A}} \overset{svn}{\approx}_\epsilon \mathbf{A}$ then $\mathbf{U}\tilde{\mathbf{A}}\mathbf{V} \overset{svn}{\approx}_\epsilon \mathbf{U}\mathbf{A}\mathbf{V}$.*

An equivalent property was not known and is in fact not true for prior notions such as unit circle, as otherwise it would show SV and UC approximation are equivalent via Item 4.

3.3 Properties of Singular Value Approximation

Here we provide several useful properties of SV approximation.

First, we show that SV approximation implies unit-circle approximation:

Lemma 3.11. *If $\mathbf{A}, \tilde{\mathbf{A}} \in \mathbb{C}^{n \times n}$ with $\tilde{\mathbf{A}} \overset{\mathbf{D}}{\approx}_\epsilon \mathbf{A}$ then $\tilde{\mathbf{A}} \overset{\circ}{\approx}_\epsilon \mathbf{A}$ with respect to degree matrix \mathbf{D} .*

Interestingly, even for 2×2 symmetric matrices, SV approximation can be separated arbitrarily from UC approximation.

Proposition 3.12. *For every $\alpha, \epsilon \in (0, 1)$, there are symmetric matrices $\mathbf{W}, \tilde{\mathbf{W}}$ with $\|\mathbf{W}\|, \|\tilde{\mathbf{W}}\| \leq 1$ such that $\tilde{\mathbf{W}} \overset{\circ}{\approx}_\epsilon \mathbf{W}$ but $\tilde{\mathbf{W}}$ is not an ϵ SV-approximation of \mathbf{W} with respect to $\mathbf{I} - \mathbf{W}$ for any $\epsilon' < \frac{\epsilon}{\sqrt{1-\alpha^2}}$.*

In prior work, ([AKM⁺20, Proposition 4.1]), it was also shown that UC approximation can be arbitrary separated from standard approximation, even for undirected graphs.

Proposition 3.13 ([AKM⁺20]). *For every $\epsilon \in (0, 1)$, there exist undirected regular graphs with random walk matrices $\tilde{\mathbf{W}}, \mathbf{W}$ such that $\tilde{\mathbf{W}}$ is an ϵ -approximation of \mathbf{W} with respect to $\mathbf{I} - \mathbf{W}$ but $\tilde{\mathbf{W}} \overset{\circ}{\approx}_c \mathbf{W}$ does not hold for every $c \in \mathbb{N}$.*

Further, since UC approximation trivially implies standard approximation (as was also argued in [AKM⁺20]) we see that SV approximation can be viewed of a strengthening of both UC and standard approximation.

A corollary of Corollary 3.10 is that SV approximation is preserved under embedding in a block matrix. In particular, it shows the stronger fact that approximation is preserved even if we use a different block structure for rows and columns (i.e. we are not embedding into principal submatrices). In contrast, unit circle approximation is only known to be preserved when the embedding pattern is a block pattern given by the directed cycle; i.e., tensoring the adjacency matrix with that of the directed cycle. In fact, the separations we prove imply unit circle approximation cannot be preserved under all block embeddings, as otherwise it would imply SV approximation by Item 3 of Lemma 3.7.

Lemma 3.14 (SV preservation under arbitrary lifting). *Let $\mathbf{A}, \tilde{\mathbf{A}} \in \mathbb{C}^{m \times n}$ be matrices such that $\tilde{\mathbf{A}} \stackrel{\mathbf{D}'_{\text{in}}, \mathbf{D}'_{\text{out}}}{\approx_\varepsilon} \mathbf{A}$. Then for all integers $i, j, k, \ell \geq 0$*

$$\begin{bmatrix} \mathbf{0}^{i \times j} & \mathbf{0}^{i \times n} & \mathbf{0}^{i \times k} \\ \mathbf{0}^{m \times j} & \tilde{\mathbf{A}} & \mathbf{0}^{m \times k} \\ \mathbf{0}^{\ell \times j} & \mathbf{0}^{\ell \times n} & \mathbf{0}^{\ell \times k} \end{bmatrix} \stackrel{\mathbf{D}'_{\text{in}}, \mathbf{D}'_{\text{out}}}{\approx_\varepsilon} \begin{bmatrix} \mathbf{0}^{i \times j} & \mathbf{0}^{i \times n} & \mathbf{0}^{i \times k} \\ \mathbf{0}^{m \times j} & \mathbf{A} & \mathbf{0}^{m \times k} \\ \mathbf{0}^{\ell \times j} & \mathbf{0}^{\ell \times n} & \mathbf{0}^{\ell \times k} \end{bmatrix}.$$

Where

$$\mathbf{D}'_{\text{in}} = \begin{bmatrix} \mathbf{0}^{i \times i} & \mathbf{0}^{i \times n} & \mathbf{0}^{i \times \ell} \\ \mathbf{0}^{n \times i} & \mathbf{D}_{\text{in}} & \mathbf{0}^{n \times \ell} \\ \mathbf{0}^{\ell \times i} & \mathbf{0}^{\ell \times n} & \mathbf{0}^{\ell \times \ell} \end{bmatrix}, \quad \mathbf{D}'_{\text{out}} = \begin{bmatrix} \mathbf{0}^{j \times j} & \mathbf{0}^{j \times m} & \mathbf{0}^{j \times k} \\ \mathbf{0}^{m \times j} & \mathbf{D}_{\text{out}} & \mathbf{0}^{m \times k} \\ \mathbf{0}^{k \times j} & \mathbf{0}^{k \times m} & \mathbf{0}^{k \times k} \end{bmatrix}.$$

Consequently, if $\tilde{\mathbf{A}} \stackrel{sv}{\approx_\varepsilon} \mathbf{A}$ then

$$\begin{bmatrix} \mathbf{0}^{i \times j} & \mathbf{0}^{i \times n} & \mathbf{0}^{i \times k} \\ \mathbf{0}^{m \times j} & \tilde{\mathbf{A}} & \mathbf{0}^{m \times k} \\ \mathbf{0}^{\ell \times j} & \mathbf{0}^{\ell \times n} & \mathbf{0}^{\ell \times k} \end{bmatrix} \stackrel{sv}{\approx_\varepsilon} \begin{bmatrix} \mathbf{0}^{i \times j} & \mathbf{0}^{i \times n} & \mathbf{0}^{i \times k} \\ \mathbf{0}^{m \times j} & \mathbf{A} & \mathbf{0}^{m \times k} \\ \mathbf{0}^{\ell \times j} & \mathbf{0}^{\ell \times n} & \mathbf{0}^{\ell \times k} \end{bmatrix}.$$

By taking sums of different such liftings, one can obtain approximation for arbitrary tensorings: if $\tilde{\mathbf{A}} \stackrel{sv}{\approx_\varepsilon} \mathbf{A}$, for any $\mathbf{M} \in \{0, 1\}^{i \times j}$, we have $\tilde{\mathbf{A}} \otimes \mathbf{M} \stackrel{sv}{\approx_\varepsilon} \mathbf{A} \otimes \mathbf{M}$. A special case of this is concatenation:

Corollary 3.15 (SV preservation under concatenation). *Let $\mathbf{A}_1, \mathbf{A}_2, \tilde{\mathbf{A}}_1, \tilde{\mathbf{A}}_2 \in \mathbb{C}^{m \times n}$ be matrices such that $\tilde{\mathbf{A}}_1 \stackrel{sv}{\approx_\varepsilon} \mathbf{A}_1$ and $\tilde{\mathbf{A}}_2 \stackrel{sv}{\approx_\varepsilon} \mathbf{A}_2$ then $\begin{bmatrix} \tilde{\mathbf{A}}_1 & \tilde{\mathbf{A}}_2 \end{bmatrix} \stackrel{sv}{\approx_\varepsilon} \begin{bmatrix} \mathbf{A}_1 & \mathbf{A}_2 \end{bmatrix}$.*

Another corollary is that SV approximation is preserved under products of different walk matrices with essentially no loss in the approximation quality. In contrast, the closest property known to be achieved by definitions of approximation considered in prior work is that unit circle approximation is preserved with no loss under only *powers* of the same walk matrix.

Lemma 3.16 (SV preservation under products). *Let $(\mathbf{N}_i)_{i \in [\ell]}, (\tilde{\mathbf{N}}_i)_{i \in [\ell]} \in \mathbb{C}^{n \times n}$ be such that for every i , $\tilde{\mathbf{N}}_i \stackrel{sv}{\approx_\varepsilon} \mathbf{N}_i$. Then $\tilde{\mathbf{N}}_\ell \cdots \tilde{\mathbf{N}}_2 \tilde{\mathbf{N}}_1 \stackrel{sv}{\approx_{\varepsilon + O(\varepsilon^2)}} \mathbf{N}_\ell \cdots \mathbf{N}_2 \mathbf{N}_1$.*

A useful application of this result is to sparsifying powers of a walk matrix of an Eulerian digraph (Theorem 4.17).

SV approximation between digraphs requires exact preservation of the degrees:

Lemma 3.17. *If $\tilde{\mathbf{A}} \stackrel{sv}{\approx_\varepsilon} \mathbf{A}$, then $\tilde{\mathbf{A}} \vec{\mathbf{1}} = \mathbf{A} \vec{\mathbf{1}}$ and $\tilde{\mathbf{A}}^\top \vec{\mathbf{1}} = \mathbf{A}^\top \vec{\mathbf{1}}$.*

SV sparsification is additive. We note that the analogous statement for normalized SV approximation is that it is preserved under convex combinations.

Lemma 3.18. *If $\tilde{\mathbf{A}}_i \stackrel{(\mathbf{D}_{\text{in}})_i, (\mathbf{D}_{\text{out}})_i}{\approx_\varepsilon} \mathbf{A}_i$ for all $i \in [k]$, letting $\mathbf{D}_{\text{in}} \stackrel{\text{def}}{=} \sum_{i \in [k]} (\mathbf{D}_{\text{in}})_i$ and $\mathbf{D}_{\text{out}} \stackrel{\text{def}}{=} \sum_{i \in [k]} (\mathbf{D}_{\text{out}})_i$, then $\sum_{i \in [k]} \tilde{\mathbf{A}}_i \stackrel{\mathbf{D}_{\text{in}}, \mathbf{D}_{\text{out}}}{\approx_\varepsilon} \sum_{i \in [k]} \mathbf{A}_i$. Consequently, if $\tilde{\mathbf{A}}_i \stackrel{sv}{\approx_\varepsilon} \mathbf{A}_i$ for every i , then $\sum_{i \in [k]} \tilde{\mathbf{A}}_i \stackrel{sv}{\approx_\varepsilon} \sum_{i \in [k]} \mathbf{A}_i$.*

SV approximation satisfies an approximate triangle inequality:

Lemma 3.19. *If $\mathbf{A}_3 \stackrel{\mathbf{D}_{\text{in}}, \mathbf{D}_{\text{out}}}{\approx_\delta} \mathbf{A}_2$ and $\mathbf{A}_2 \stackrel{\mathbf{D}_{\text{in}}, \mathbf{D}_{\text{out}}}{\approx_\varepsilon} \mathbf{A}_1$ then $\mathbf{A}_3 \stackrel{\mathbf{D}_{\text{in}}, \mathbf{D}_{\text{out}}}{\approx_{\varepsilon+\delta+\varepsilon\delta}} \mathbf{A}_1$. Consequently, if $\mathbf{A}_3 \stackrel{sv}{\approx_\delta} \mathbf{A}_2$ and $\mathbf{A}_2 \stackrel{sv}{\approx_\varepsilon} \mathbf{A}_1$ then $\mathbf{A}_3 \stackrel{sv}{\approx_{\varepsilon+\delta+\varepsilon\delta}} \mathbf{A}_1$, and if $\mathbf{A}_3 \stackrel{sun}{\approx_\delta} \mathbf{A}_2$ and $\mathbf{A}_2 \stackrel{sun}{\approx_\varepsilon} \mathbf{A}_1$ then $\mathbf{A}_3 \stackrel{sun}{\approx_{\varepsilon+\delta+\varepsilon\delta}} \mathbf{A}_1$. Moreover, if for $\delta \in (0, 1/2)$ and $\mathbf{A}_0, \dots, \mathbf{A}_\ell$ we have $\mathbf{A}_i \stackrel{sv}{\approx_{\delta/2^\ell}} \mathbf{A}_{i-1}$ for every i , then $\mathbf{A}_\ell \stackrel{sv}{\approx_\delta} \mathbf{A}_0$. Moreover, the equivalent claim holds for normalized SV approximation.*

A regular undirected graph SV approximates the complete graph with error equal to its expansion.

Lemma 3.20. *Let G be a strongly connected, d -regular directed multigraph on n vertices with adjacency matrix \mathbf{A} and let $\mathbf{J} \in \mathbb{R}^{n \times n}$ be a matrix with $1/n$ in every entry (i.e. \mathbf{J} is the walk matrix of the complete graph with a self loop on every vertex). Then $\lambda(G) \leq 1 - \lambda/2$ if and only if $\mathbf{A}/d \stackrel{sun}{\approx_\lambda} \mathbf{J}$.*

4 Sparsification

Given the Laplacian of an Eulerian directed graph, we wish to compute a sparsifier with respect to SV approximation. In this section, we show how to solve a more general problem of sparsifying a non-negative rectangular matrix $\mathbf{A} \in \mathbb{R}_{\geq 0}^{m \times n}$ with respect to SV approximation. Further, in Section 4.5 we show how to construct square sparsifiers via the derandomized square, and in Section 4.6 we show how to sparsify powers of walk matrices. We first give an informal overview of our approach.

We give a series of reductions that reduce this problem to degree- and bipartition-preserving sparsification of expander graphs with respect to ε -approximation. By Lemma 3.7 (Item 2), to SV-sparsify (possibly directed) \mathbf{A} , it suffices to SV-sparsify the undirected bipartite lift

$$\begin{bmatrix} \mathbf{0} & \mathbf{A}^\top \\ \mathbf{A} & \mathbf{0} \end{bmatrix}$$

with respect to SV approximation. In particular, given a symmetric matrix satisfying $\text{slift}(\tilde{\mathbf{A}}) \stackrel{sv}{\approx_\varepsilon} \text{slift}(\mathbf{A})$, we have by Lemma 3.7 that $\tilde{\mathbf{A}} \stackrel{sv}{\approx_\varepsilon} \mathbf{A}$. Thus, any symmetric bipartition-preserving SV sparsifier of undirected graphs immediately gives an SV sparsifier for Eulerian digraphs. In fact, every SV sparsifier of undirected graphs must be bipartition-preserving, and an asymmetric sparsifier can be used to read off sparsifiers of the underlying directed graphs. For a formal statement of this stronger claim, see Lemma A.5.

We show (Lemma 4.3) that for undirected bipartite *expanders*, degree- and bipartition-preserving approximation with respect to \mathbf{D} implies SV approximation.

From this, we can construct SV sparsifiers (Theorem 4.11) by computing an expander decomposition of the bipartite lift, then sparsifying the relevant subgraphs in a suitable way. While we can use existing techniques to obtain this (with worse log factors), we also develop a new sparsification approach based on cycle decompositions. We show that in a bipartite expander, decomposing edges incident to high-degree vertices into edge-disjoint cycles, then taking the odd or even edges

of each cycle with probability $1/2$ produces a sparsifier with high probability. This sparsification procedure automatically preserves degrees and bipartitions, which we require to lift approximation with respect to \mathbf{D} to SV approximation. Prior work [CGP⁺18] showed an equivalent result in general Eulerian digraphs, except they required the cycles to be short, resulting in almost-linear runtime [CGP⁺18, PY19].

For convenience, we define notation for the adjacency and degree matrices of graphs:

Definition 4.1. For the remainder of the section, for a weighted undirected graph G we let \mathbf{A}_G denote the adjacency matrix of G and $\mathbf{D}_G \stackrel{\text{def}}{=} \mathbf{A}_G \vec{\mathbf{1}}$ denote the degree matrix. As such, an ε -SV sparsifier of G is a graph H such that $\mathbf{A}_H \stackrel{sv}{\approx}_\varepsilon \mathbf{A}_G$.

First, we show that for undirected *expanders*, degree- and bipartition-preserving sparsification with respect to the degree matrix implies sparsification with respect to SV approximation.

Definition 4.2 (Bipartiteness). We say $\mathbf{A} \in \mathbb{R}_{\geq 0}^{n \times n}$ is *bipartite* with *bipartition* $S, T \subseteq [n]$ if S and T partition V and $\mathbf{A}_{S,S} = \mathbf{0}_{S \times S}$ and $\mathbf{A}_{T,T} = \mathbf{0}_{T \times T}$. Given a graph G with adjacency matrix \mathbf{A} , we say that a sparsifier H is *bipartition-preserving* if for every bipartition S, T in G , we have $H_{S,S} = \mathbf{0}_{S \times S}$ and $H_{T,T} = \mathbf{0}_{T \times T}$ (where the statement is vacuously satisfied if G is not bipartite).

In the following lemma we show that for non-negative matrices that are bipartite with the same bipartition, SV approximation is implied by approximation with respect to the error matrix \mathbf{D} (Definition 3.1), up to a loss in the condition number.

Lemma 4.3 (From diagonal approximation to SV approximation). *Suppose that symmetric $\tilde{\mathbf{A}} \in \mathbb{R}_{\geq 0}^{n \times n}$ is a degree and bipartition-preserving $\varepsilon/2$ -approximation of symmetric bipartite $\mathbf{A} \in \mathbb{R}_{\geq 0}^{n \times n}$ with respect to error matrix $\mathbf{D} \stackrel{\text{def}}{=} \text{diag}(\mathbf{A}\vec{\mathbf{1}}) = \text{diag}(\mathbf{A}^\top \vec{\mathbf{1}})$. Then $\tilde{\mathbf{A}} \stackrel{sv}{\approx}_\gamma \mathbf{A}$ for $\gamma \stackrel{\text{def}}{=} \frac{\varepsilon}{\lambda^2}$ where $\lambda \stackrel{\text{def}}{=} 1 - \lambda_2(\mathbf{D}^{+1/2} \mathbf{A} \mathbf{D}^{+1/2})$.*

Proof. Recall by Lemma 3.7 that it suffices to show that $\tilde{\mathbf{A}}$ is an $\varepsilon/2$ -approximation of \mathbf{A} with respect to $\mathbf{E} \stackrel{\text{def}}{=} \mathbf{D} - \mathbf{A} \mathbf{D}^+ \mathbf{A}$. First, we have

$$\mathbf{E} \succeq (\mathbf{D} - \mathbf{A}) \mathbf{D}^+ (\mathbf{D} + \mathbf{A}) \succeq \mathbf{0}.$$

Furthermore by Lemma 3.2 and the assumption that $\tilde{\mathbf{A}}$ is an $\varepsilon/2$ -approximation of \mathbf{A} with respect to \mathbf{D} we have

$$-\varepsilon/2 \cdot \mathbf{D} \preceq \mathbf{A} - \tilde{\mathbf{A}} \preceq \varepsilon/2 \cdot \mathbf{D}.$$

By assumption that \mathbf{A} has expansion λ and is bipartite (and hence its spectrum is symmetric), we have

$$(\mathbf{D} - \mathbf{A})^+ \preceq \frac{1}{\lambda} \mathbf{D}^+, \quad (\mathbf{D} + \mathbf{A})^+ \preceq \frac{1}{\lambda} \mathbf{D}^+.$$

Furthermore, as \mathbf{A} is the adjacency matrix of an undirected graph the only singular values of \mathbf{A} of magnitude 1 can be -1 and 1 . Thus since $\tilde{\mathbf{A}}$ matches degrees exactly and preserves bipartitions we have that $\ker(\mathbf{A} - \tilde{\mathbf{A}}) \supseteq \ker(\mathbf{E})$. Thus we have

$$\begin{aligned} \left\| \mathbf{E}^{+1/2} (\mathbf{A} - \tilde{\mathbf{A}}) \mathbf{E}^{+1/2} \right\| &\leq \left\| (\mathbf{D} + \mathbf{A})^{+1/2} \mathbf{D}^{1/2} (\mathbf{D} - \mathbf{A})^{+1/2} (\mathbf{A} - \tilde{\mathbf{A}}) (\mathbf{D} - \mathbf{A})^{+1/2} \mathbf{D}^{1/2} (\mathbf{D} + \mathbf{A})^{+1/2} \right\| \\ &\leq \frac{1}{\lambda} \left\| \mathbf{D}^{+1/2} \mathbf{D}^{1/2} (\mathbf{D} - \mathbf{A})^{+1/2} (\mathbf{A} - \tilde{\mathbf{A}}) (\mathbf{D} - \mathbf{A})^{+1/2} \mathbf{D}^{1/2} \mathbf{D}^{+1/2} \right\| \\ &\leq \frac{1}{\lambda} \left\| (\mathbf{D} - \mathbf{A})^{+1/2} (\mathbf{A} - \tilde{\mathbf{A}}) (\mathbf{D} - \mathbf{A})^{+1/2} \right\| \\ &\leq \frac{1}{\lambda^2} \left\| \mathbf{D}^{+1/2} (\mathbf{A} - \tilde{\mathbf{A}}) \mathbf{D}^{+1/2} \right\| \leq \frac{\varepsilon}{\lambda^2}. \quad \square \end{aligned}$$

4.1 Cycle and Expander Decompositions

We give a new analysis of degree-preserving expander sparsification.

Chu, Gao, Peng, Sachdeva, Sawlani, and Wang [CGP⁺18] constructed sparsifiers for Eulerian digraphs, which we now discuss. They decomposed the low-importance edges of a graph into a series of edge-disjoint cycles C_1, \dots, C_t . Then for each cycle, they chose a random orientation (clockwise or counterclockwise) and kept every edge in this orientation, and upweighted these by a factor of two. One can see that this procedure gives a subgraph and maintains Eulerianness. They bounded the impact of a single such sample in terms of the length of the cycle, and by giving a sophisticated algorithm for decomposing a graph into many short cycles concluded an almost-linear time sparsifier for standard approximation.

We follow this approach but make two modifications. First, we require the cycles to be *Forward-Backward (FB)*, where we always alternate between clockwise and counterclockwise edges (and all cycles are of even length). This ensures that we exactly preserve *degrees*, not just Eulerianness, which is essential for SV approximation. We take advantage of the reduction from directed to undirected SV sparsification to perform this decomposition in the undirected bipartite lift, where undirected cycles correspond to alternating cycles in the original graph. Our second modification is to prove a bound on the importance of the cycles that is independent of their length, as long as the cycles lie inside an expander. This enables us to use a simple linear time algorithm for finding the decomposition.

We first recall the standard procedure that enables us to efficiently decompose an undirected graph into a union of (potentially long) FB cycles.

Lemma 4.4. *Given an unweighted undirected graph $G = (V, E)$, the algorithm CYCLEDECOMP returns a collection of edge-disjoint cycles $C_1, \dots, C_T \subseteq E$ in time $O(|E| + |V|)$ such that at most n edges are not contained in some cycle.*

Algorithm 1: CYCLEDECOMP ($G = (V, E)$), G undirected

```

1 Initialize  $E_{ex}, \mathcal{C} = \{\}$ .
2 for  $v \in V$  do
3   while Greedily construct a non-backtracking path  $v = u_0, u_1, \dots, u_k$  in an arbitrary
   fashion and mark visited vertices. do
4     if  $u_k$  has no neighbors other than  $u_{k-1}$  then
5       Add  $E_{ex} \leftarrow E_{ex} \cup \{(u_{k-1}, u_k)\}$  and remove  $(u_{k-1}, u_k)$  from  $E$ .
6       Continue search from  $u_{k-1}$ .
7     end
8     if  $u_k = u_{k'}$  for some  $k' < k - 1$  then
9       Let  $\mathcal{C} \leftarrow \mathcal{C} \cup (u_{k'}, u_{k'+1}, \dots, u_{k-1})$  and let
        $E \leftarrow E \setminus \{(u_{k'}, u_{k'+1}), (u_{k'+1}, u_{k'+2}), \dots, (u_{k-1}, u_k)\}$ .
10      Continue search from  $u_k$ .
11    end
12  end
13 end
14 return  $\mathcal{C}, E_{ex}$ 

```

Proof. The runtime and the fact that the cycles are edge disjoint is direct from the description.

Then note that for every edge (u, v) that we add to E_{ex} (and remove from E) we must create an isolated vertex v . But this can happen at most n times, so we are done. \square

Furthermore, we recall the definition of an expander partition:

Definition 4.5. Given ϕ, γ and an undirected graph G , a (ϕ, γ) -*expander partition* of G is a partition V_1, \dots, V_t such that at most $\gamma \cdot m$ edges cross between partitions and for every $i \in [t]$, $\lambda(G[V_i]) \leq 1 - \phi$.

We will use algorithms for expander decomposition as a key subroutine in our sparsifier. We require a small constant fraction of edges to lie between clusters, and thus state the algorithms fixing $\gamma = 1/16$ such that $m/16$ edges cross subgraphs. In particular, given an algorithm `EXPANDERDECOMP` computing a $(\Phi, m/16)$ expander decomposition, let $T(\text{EXPANDERDECOMP}, m)$ be its runtime (on graphs with m edges) and $\Phi(\text{EXPANDERDECOMP}) = \Phi$. We recall the best known nearly-linear expander partitioning algorithm, as well as the (optimal) existential result:

Theorem 4.6 ([ADK22]). *There is an algorithm `EXPANDERDECOMP`(G) running in randomized time $T(\text{EXPANDERDECOMP}, m) = O(m \log^7 m)$ that computes a $(\Phi = \log^{-4} m, 1/16)$ expander decomposition of G with high probability.*

Theorem 4.7 ([ST04]). *There exists a $(\Phi = \log^{-2} m, 1/16)$ expander decomposition of G .*

4.2 Sparsifying Bipartite Expanders

The core of our algorithm is a way to sparsify bipartite, unweighted expanders in a degree- and bipartition-preserving way.

The algorithm decomposes the graph into a series of edge-disjoint cycles, then takes either twice the odd or twice the even edges with probability $1/2$. In the case that the undirected graph corresponds to the bipartite lift of an Eulerian digraph G , this precisely corresponds to decomposing G into a union of Forward-Backward cycles and taking the forward or backward edges for each cycle. We bound the error in terms of the degree matrix \mathbf{D} . This is tolerable as we apply this procedure inside expanders, so by Lemma 4.3 we can transform this into an SV approximation guarantee with $\text{polylog}(n)$ loss in approximation quality.

Lemma 4.8. *There is a constant $c > 0$ such that given $\delta \in (0, 1)$ and an unweighted, undirected bipartite graph G with m edges, `SPARSIFYCYCLEUNWEIGHTED` returns in time $O(m)$ a graph H with edge weights in $\{1, 2\}$ such that with high probability \mathbf{A}_H is a δ -approximation of \mathbf{A}_G with respect to error matrix \mathbf{D}_G , and H has at most $\max\{cn\delta^{-2} \log n, (7/8)m\}$ edges and exactly preserves degrees and bipartiteness.*

To prove correctness, we recall a result on matrix concentration of Tropp:

Lemma 4.9 ([Tro12]). *Let $\{\mathbf{X}_i\}$ be $d \times d$ random PSD matrices such that $\mathbb{E}[\sum_i \mathbf{X}_i] \preceq \mathbf{X}$. For $\delta \leq 1$,*

$$\Pr \left[\sum_i \mathbf{X}_i - \mathbb{E} \left[\sum_i \mathbf{X}_i \right] \succeq \delta \mathbf{X} \right] \leq 1 - d \cdot e^{-\delta^2/3R}, \quad \Pr \left[\sum_i \mathbf{X}_i - \mathbb{E} \left[\sum_i \mathbf{X}_i \right] \preceq -\delta \mathbf{X} \right] \leq 1 - d \cdot e^{-\delta^2/3R},$$

provided that $R \geq \|\mathbf{X}^{+1/2} \mathbf{X}_i \mathbf{X}^{+1/2}\|$ almost surely.

Using this matrix concentration result, we can prove the theorem.

Algorithm 2: SPARSIFYCYCLEUNWEIGHTED $(\delta, G = (V, E))$

```

1 if  $\delta < \sqrt{cn \log n/m}$  then
2   | return  $G$ 
3 end
4 Let  $d_{low} \stackrel{\text{def}}{=} \frac{m}{2n}$ .
5 Let  $E_{low}$  be the edges adjacent to vertices  $v$  with  $d(v) \leq d_{low}$ .
6 Let  $E_{ex}, (C_1, \dots, C_t) = \text{CYCLEDECOMP}(V, E \setminus E_{low})$ .
7 Initialize  $H = (V, E_{ex} \cup E \setminus E_{low})$ .
8 for  $i = 1, \dots, t$  do
9   | Let  $(a_1, b_1), (b_1, a_2), \dots, (b_k, a_1) = C_i$ .
10  | Choose  $b \leftarrow 0, 1$  uniformly at random.
11  | if  $b = 0$  then
12  |   |  $H \leftarrow H \cup \{2 \cdot (a_i, b_i)\}_{i \in [k]}$  // Add with weight 2
13  |   end
14  | else  $H \leftarrow H \cup \{2 \cdot (b_i, a_{i+1})\}_{i \in [k]}$ ;
15 end
16 return  $H$ 

```

Proof of Lemma 4.8. First, note that if we do not sparsify G at all we have $cn\delta^{-2} \log n < m$ and hence the sparsity requirement is already satisfied. Subsequently we assume this is not the case.

Recall E_{low} is the set of edges adjacent to a vertex with degree at most $d_{low} = (m/2n)$, and observe that there are at most $d_{low} \cdot n \leq m/2$ such edges. Further recall C_1, \dots, C_t are the cycles found by CYCLEDECOMP applied to $E \setminus E_{low}$. By assumption that G is bipartite, all such cycles are of even length.

Let $\mathbf{X}_i \stackrel{\text{def}}{=} \tilde{\mathbf{D}}_i - \tilde{\mathbf{A}}_i$ be the random variable of the Laplacian of the i th sampled cycle and observe $\mathbf{0} \preceq \mathbf{X}_i \preceq 2\mathbf{I}_{C_i}$. Furthermore, let $\mathbf{X} \stackrel{\text{def}}{=} \mathbf{D}_G \succeq \mathbf{0}$. We have

$$\mathbb{E} \left[\sum_i \mathbf{X}_i \right] = \sum_{i=1}^t \mathbf{L}_{C_i} \preceq \mathbf{X}.$$

Furthermore, for every $i \in [t]$ we have

$$\left\| \mathbf{X}^{+/2} \mathbf{X}_i \mathbf{X}^{+/2} \right\| \leq 2 \left\| \mathbf{D}^{+/2} \mathbf{I}_{C_i} \mathbf{D}^{+/2} \right\| \leq \frac{4n}{m}$$

where the final step follows from all cycles being exclusively incident to vertices with degree at least $m/2n$. Thus, we apply Lemma 4.9 with $\{\mathbf{X}_i\}_{i \in [t]}$, $R = 4n/m$, $\mathbf{X} = \mathbf{X}$ and $\delta = \delta/c$. Letting $\mathbf{D}_C - \mathbf{A}_C$ be the random graph obtained by the cycle sampling procedure, with high probability we have

$$-\delta \mathbf{D}_G \preceq (\mathbf{D}_C - \mathbf{A}_C) - \sum_{i=1}^t (\mathbf{D}_{C_i} - \mathbf{A}_{C_i}) \preceq \delta \mathbf{D}_G.$$

By adding the non-random components to we obtain

$$-\delta \mathbf{D}_G \preceq \mathbf{A}_H - \mathbf{A}_G \preceq \delta \mathbf{D}_G$$

and thus \mathbf{A}_H is an δ -approximation of \mathbf{A}_G with respect to \mathbf{D}_G with high probability.

Furthermore, by Lemma 4.4 at least $m/2 - n \geq m/4$ edges are included in the cycles C_1, \dots, C_t , and hence H has at most $7m/8$ edges under the assumption that $m \geq cn$, and H preserves degrees and bipartiteness by construction. \square

4.3 Sparsifying a Constant Fraction of Edges

We then construct our routine for sparsifying a graph with respect to SV approximation. At each iteration, we decrease the number of edges by a constant factor. We first bucket edges into powers of two, then apply SPARSIFYCYCLEUNWEIGHTED.

Lemma 4.10. *Given $\varepsilon > 0$ and $b \in \mathbb{N}$ and an undirected bipartite graph G with m edges, each with weight in $\{1, 2, 2^2, \dots, 2^b\}$, SPARSIFYCYCLE returns a graph H with*

$$O\left(b \cdot n\varepsilon^{-2}\Phi^{-4} \log n\right) + \frac{15}{16}m$$

edges where $\Phi = \Phi(\text{EXPANDERDECOMP})$, each with weight in $\{1, 2, 2^2, \dots, 2^{b+1}\}$, such that with high probability $\mathbf{A}_H \stackrel{sv}{\approx}_\varepsilon \mathbf{A}_G$. Moreover, SPARSIFYCYCLE runs in randomized time $O(m + bn + \mathbf{T}(\text{EXPANDERDECOMP}, m))$.

Algorithm 3: SPARSIFYCYCLE ($\varepsilon, b, G = (V, E)$)

```

1 Let  $\delta = \varepsilon^{-1}\Phi^2$ .
2 Initialize  $H = (V, \emptyset)$ .
3 for  $i = 0, \dots, b$  do
4   Initialize (unweighted)  $G_i$  as  $G$  restricted to edges with weight  $2^i$ .
5   Let  $V_{i,1}, \dots, V_{i,t}, E_{ex} \stackrel{\text{def}}{=} \text{EXPANDERDECOMP}(G_i)$ .
6   Let  $H \leftarrow H \cup 2^i \cdot E_{ex}$ .
7   for  $j \in 1, \dots, t$  do
8     Let  $H_{i,j} \stackrel{\text{def}}{=} \text{SPARSIFYCYCLEUNWEIGHTED}(\delta, G_i[V_{i,j}])$ 
9     Let  $H \leftarrow H \cup 2^i \cdot H_{i,j}$ .
10  end
11 end
12 return  $H$ 

```

Proof. Let G_0, \dots, G_b be the unweighted bipartite graphs where G_i contains an edge (u, v) if there is an equivalent edge with weight 2^i . Thus, observe that

$$\sum_{i=0}^b 2^i \cdot (\mathbf{D}_{G_i} - \mathbf{A}_{G_i}) = \mathbf{D}_G - \mathbf{A}_G$$

and it suffices to sparsify unweighted bipartite graphs. Furthermore let $m_i \stackrel{\text{def}}{=} |E(G_i)|$ and observe $\sum_i m_i = m$.

Next, for every i observe that EXPANDERDECOMP computes a decomposition $V_{i,1}, \dots, V_{i,t}$ such that letting $G_{i,j} \stackrel{\text{def}}{=} G_i[V_{i,j}]$ we have $\lambda(G_{i,j}) \leq 1 - \Phi$, and at most $m_i/16$ edges do not lie in one of these subgraphs. Let $n_{i,j} \stackrel{\text{def}}{=} |G_{i,j}|$ and observe $\sum_j n_{i,j} \leq n$ and let $m_{i,j} \stackrel{\text{def}}{=} |E(G_{i,j})|$ and observe $\sum_j m_{i,j} \leq m_i$. By Lemma 4.8, we have that for every i, j , $\mathbf{A}_{H_{i,j}}$ is a degree and bipartition-preserving δ -approximation of $\mathbf{A}_{G_{i,j}}$ with respect to $\mathbf{D}_{G_{i,j}}$ with high probability, and thus by Lemma 4.3

$$\mathbf{A}_{H_{i,j}} \stackrel{sv}{\approx}_{\delta \cdot \Phi^{-2}} \mathbf{A}_{G_{i,j}}.$$

Thus by Lemma 3.18 we have

$$\mathbf{A}_H \stackrel{sv}{\approx}_{\delta \cdot \Phi^{-2}} \mathbf{A}_G$$

and so by our choice of $\delta = \varepsilon \Phi^2$ we obtain the desired approximation.

We now analyze the number of edges. The number of edges not placed into a subgraph $G_{i,j}$ is at most

$$\sum_{i=1}^b \frac{m_i}{16} = \frac{m}{16}.$$

By Lemma 4.8 the number of edges in the sparsified graphs is at most

$$\begin{aligned} \sum_{i=1}^b \sum_{j=1}^t \max\{cn_{i,j}\delta^{-2} \log n, \frac{7}{8}m_{i,j}\} &\leq \sum_{i=1}^b \sum_{j=1}^t \left(cn_{i,j}\delta^{-2} \log n + \frac{7}{8}m_{i,j} \right) \\ &= O(bn\varepsilon^{-2}\Phi^{-4} \log n) + \frac{7}{8} \sum_{i=1}^b \sum_{j=1}^t m_{i,j} \\ &= O(bn\varepsilon^{-2}\Phi^{-4} \log n) + \frac{7}{8}m \end{aligned}$$

and hence the total number of edges is bounded as desired. Finally, we analyze the time. We can compute every G_i with a single pass through the edge list, so the total work is bounded by $\sum_i \sum_j O(m_{i,j} + n_{i,j}) + \sum_{i=1}^b \mathsf{T}(\text{EXPANDERDECOMP}, m_i) = O(m + bn + \mathsf{T}(\text{EXPANDERDECOMP}, m))$ as claimed. \square

4.4 Nearly Linear-Sized Sparsifiers

We now apply our routine for sparsifying a constant fraction of edges in the natural recursive way.

Theorem 4.11. *Given $\varepsilon \in (0, 1)$ and an undirected bipartite graph G with m edges with integer edge weights in $[1, U]$, SPARSIFYGRAPH returns a graph H with*

$$O(\log(nU) \cdot n\varepsilon^{-2}\Phi^{-4} \log^3 n)$$

edges where $\Phi = \Phi(\text{EXPANDERDECOMP})$, such that with high probability $\mathbf{A}_H \stackrel{sv}{\approx}_{\varepsilon} \mathbf{A}_G$. Moreover, SPARSIFYGRAPH runs in time $O(m + nb + \mathsf{T}(\text{EXPANDERDECOMP}, m \log(U)))$.

Proof. First observe that in Phase 1 of the algorithm, we obtain H_0 such that H_0 has at most mb edges (recalling $b = \lceil \log(U) \rceil$) and each edge has weight $\{1, 2, 2^2, \dots, 2^b\}$, and $\mathbf{A}_{H_0} = \mathbf{A}_G$. Then in each phase we obtain H_j such that

$$\mathbf{A}_{H_j} \stackrel{sv}{\approx}_{\varepsilon/2 \log n} \mathbf{A}_{H_{j-1}}$$

with high probability by Lemma 4.10. Then applying Lemma 3.19 we obtain

$$\mathbf{A}_{H_t} \stackrel{sv}{\approx}_{\varepsilon} \mathbf{A}_G$$

with high probability as desired. We next analyze the sparsity. Observe that the final edge weights are bounded by $2^{b+t} = \text{poly}(nU)$ and hence the final sparsity is

$$O((b+t)n\varepsilon^{-2}\Phi^{-4} \log^3 n) = O(\log(nU)n\varepsilon^{-2}\Phi^{-4} \log^3 n).$$

Algorithm 4: SPARSIFYGRAPH $(\varepsilon, G = (V, E))$

```

1 Initialize  $H_0 = (V, \emptyset)$ .
2 Let  $b \stackrel{\text{def}}{=} \lceil \log(U) \rceil$ .
3 for  $i = 0, \dots, b$  do
4   for  $(u, v) \in E$  do
5     Let  $\langle w_G(u, v) \rangle$  be the binary expansion of  $w_G(u, v)$ .
6     if  $\langle w_G(u, v) \rangle_i = 1$  then Add  $(u, v)$  to  $H_0$  with weight  $2^i$ .;
7   end
8 end
9 for  $j = 1, \dots, t \stackrel{\text{def}}{=} O(\log n)$  do
10  | Let  $H_j \leftarrow \text{SPARSIFYCYCLE}(\varepsilon/2 \log n, b + j, H_{j-1})$ .
11 end
12 return  $H_t$ 

```

Finally, we analyze the time complexity. Our initial call has mb edges and thus runs in time $O(mb + bn + \mathbf{T}(\text{EXPANDERDECOMP}, mb))$ by Lemma 4.10, and all subsequent calls are to sparsify graphs with a constant factor fewer edges, so the time is dominated by the initial term. \square

By plugging in the algorithmic (Theorem 4.6) and existential (Theorem 4.6) results for expander partitions, we obtain our sparsifier routines:

Corollary 4.12. *Given $\varepsilon \in (0, 1)$ and an undirected bipartite graph G with m edges with integer edge weights in $[1, U]$, there is a randomized algorithm that returns a graph H with*

$$O(\log(nU) \cdot n\varepsilon^{-2} \log^{19} n)$$

edges such that with high probability $\mathbf{A}_H \stackrel{sv}{\approx}_\varepsilon \mathbf{A}_G$. Moreover, the algorithm runs in time $O(\log(U)m \log^7 m)$.

Corollary 4.13. *Given $\varepsilon \in (0, 1)$ and an undirected bipartite graph G with m edges with integer edge weights in $[1, U]$, there is a graph H with*

$$O(\log(nU) \cdot n\varepsilon^{-2} \log^{11} n)$$

edges such that $\mathbf{A}_H \stackrel{sv}{\approx}_\varepsilon \mathbf{A}_G$.

4.5 Derandomized Square Sparsification

Here we show that the derandomized square of Rozenman and Vadhan [RV05] gives an SV sparsification of the square graph. We first explain the derandomized square. Given a d -regular graph G , the squaring operation can be thought of as placing a $d \times d$ complete graph between the in- and out-neighbors of every vertex v . The derandomized square replaces this complete graph with an expander. As an expander approximates the complete graph, this sparsification operation produces an approximate square.

We first formally define the derandomized square.

Definition 4.14. Given a d -regular graph $G = ([n], E)$ with neighbor function $\Gamma_G : [n] \times [d] \rightarrow [n]$ and c -regular graph $H = ([d], E')$ with neighbor function $\Gamma_H : [d] \times [c] \rightarrow [d]$, the *derandomized square of G with respect to H* , denoted $G \circledast H$, is the dc -regular graph on n vertices with neighbor function

$$\Gamma_{G \circledast H}(v, (i, j)) = \Gamma_G(\Gamma_G(v, i), \Gamma_H(i, j)).$$

Prior analyses of the derandomized square have shown that it improves expansion approximately as well as the true square [RV05] and that it produces unit-circle square sparsifiers [AKM⁺20]. We strengthen the latter conclusion to SV square sparsifiers.

Lemma 4.15. *Let $G = (V, E)$ be a d -regular directed multigraph with random walk matrix \mathbf{W} . Let H be a c -regular expander on d vertices with $\lambda(H) \leq 1 - \epsilon$ and let $\widetilde{\mathbf{W}}$ be the adjacency matrix of $G \circledast H$. Then*

$$\widetilde{\mathbf{W}} \stackrel{svn}{\approx}_\epsilon \mathbf{W}^2.$$

Proof. Let \mathbf{W}_H be the random walk matrix corresponding to H , and $\mathbf{J} \in \mathbb{R}^{d \times d}$ be the random walk matrix of a complete graph with self loops. By Lemmas 3.14 and 3.20 we have

$$\mathbf{I}_n \otimes \mathbf{W}_H \stackrel{sv}{\approx}_\epsilon \mathbf{I}_n \otimes \mathbf{J}. \quad (18)$$

Let \mathbf{F} be an $nd \times nd$ edge-rotation permutation matrix such that $\mathbf{F}((v, j), (u, i)) = 1$ if the i th edge leaving u is the j th edge entering v . It is straight-forward to verify that

$$\mathbf{W}^2 = \mathbf{Q}^\top \mathbf{F}(\mathbf{I} \otimes \mathbf{J})\mathbf{F}\mathbf{Q}$$

and

$$\widetilde{\mathbf{W}} = \mathbf{Q}^\top \mathbf{F}(\mathbf{I} \otimes \mathbf{W}_H)\mathbf{F}\mathbf{Q}$$

where $\mathbf{Q} \stackrel{\text{def}}{=} \mathbf{I} \otimes \frac{1_d}{\sqrt{d}}$ is an $nd \times n$ matrix. Now note that $\|\mathbf{F}\| \leq 1$ and $\|\mathbf{Q}\| \leq 1$ and thus $\|\mathbf{F}\mathbf{Q}\|, \|\mathbf{Q}^\top \mathbf{F}\| \leq 1$. Thus by Theorem 3.9, and Equation (18) we have

$$\widetilde{\mathbf{W}} \stackrel{sv}{\approx}_\epsilon \mathbf{W}^2. \quad \square$$

4.6 Sparsification of Random Walk Polynomials

We give the first nearly-linear time algorithm for sparsifying random walk polynomials of Eulerian digraphs with second singular value bounded away from 1. In particular, given an Eulerian digraph $\mathbf{W} \in \mathbb{R}_{\geq 0}^{n \times n}$ with second singular value bounded away from 1, we compute a sparse graph such that the walk matrix $\widetilde{\mathbf{W}}$ approximates \mathbf{W}^ℓ with respect to SV approximation. For this result, we crucially require that SV approximation is preserved under products, a feature that is not obtained by prior notions of approximation.

Definition 4.16. Given a strongly connected Eulerian digraph G , we let $\sigma(G) = \sigma_2(\mathbf{D}_G^{-1/2} \mathbf{A}_G \mathbf{D}_G^{-1/2})$ and refer to this as the *second normalized singular value* of G .

Theorem 4.17. *Given $\ell \in \mathbb{N}$ and $\epsilon \in (1/\text{poly}(n), 1)$ and an Eulerian digraph G where $\sigma(G) \leq 1 - 1/\text{poly}(n)$ with m edges with integer edge weights in $[1, \text{poly}(n)]$, SPARSIFYPOWER returns in time $\tilde{O}(m + n\epsilon^{-2} \log^3(\ell))$ an Eulerian digraph H with $\tilde{O}(n\epsilon^{-2} \log^2(\ell))$ edges with integer edge weights in $[1, \text{poly}(n, \ell, 1/\epsilon)]$ such that*

$$\mathbf{D}_H^{+1/2} \mathbf{A}_H \mathbf{D}_H^{+1/2} \stackrel{svn}{\approx}_\epsilon (\mathbf{D}_G^{+1/2} \mathbf{A}_G \mathbf{D}_G^{+1/2})^\ell.$$

This implies the following corollary.

Definition 4.18. Let \mathbf{A} be the adjacency matrix of an Eulerian digraph with diagonal matrix of degrees \mathbf{D} and walk matrix \mathbf{W} . The k th random walk power of \mathbf{A} is defined as

$$\mathbf{D}^{1/2} (\mathbf{D}^{-1/2} \mathbf{A} \mathbf{D}^{-1/2})^k \mathbf{D}^{1/2} = \mathbf{W}^k \mathbf{D}.$$

Corollary 4.19. *Let G be an Eulerian digraph with $\text{poly}(n)$ weak mixing time, adjacency matrix \mathbf{A} , diagonal degree matrix \mathbf{D} , and walk matrix \mathbf{W} . Then for any positive integer $k \leq \text{poly}(n)$ and $\gamma \geq 1/\text{poly}(n)$, we can, in nearly linear time, output an $\tilde{O}(n/\varepsilon^2)$ sized ε -SV-sparsifier of k th random walk power of $\mathbf{A}' = (1 - \gamma)\mathbf{A} + \gamma\mathbf{D}$. Furthermore, the walk matrix of the true k th random walk power of \mathbf{A}' is given by $((1 - \gamma)\mathbf{W} + \gamma\mathbf{I})^k$ so it is the k step γ -lazy random walk on G .*

Proof. Immediate from Lemma B.5 and Theorem 4.17. \square

To prove these, we first recall the square sparsification algorithm implicit in the work of Cohen et al. [CKP⁺17].

Theorem 4.20 (Implicit in [CKP⁺17]). *Given an Eulerian weighted digraph G with Laplacian $\mathbf{D} - \mathbf{A}_G$, let G^2 be the graph with Laplacian $\mathbf{D} - \mathbf{A}_G\mathbf{D} + \mathbf{A}_G$. Let \mathbf{U} denote the Laplacian of the undirected graph corresponding to $(\mathbf{A}_{G^2} + \mathbf{A}_{G^2}^T)/2$. Then SPARSIFY SQUARE of [CKP⁺17] produces in time $\tilde{O}(m)$, a weighted graph H with the following properties:*

1. H has the same weighted in- and out- degree sequences as those of G^2 .
2. H has $\tilde{O}(m/\varepsilon^2)$ nonzero edges.
3. \mathbf{A}_H is an $\varepsilon/2$ -approximation of $\mathbf{A}_G\mathbf{D}^{-1}\mathbf{A}_G$ with respect to $\mathbf{S}_{\mathbf{D} - \mathbf{A}_G\mathbf{D} + \mathbf{A}_G}$.

We can use a block embedding and Theorem 4.20 to obtain a nearly-linear time algorithm for sparsifying products with respect to SV approximation.

Lemma 4.21. *Given $\varepsilon > 0$ and Eulerian digraphs G_1, G_2 each with at most m edges with matching degree matrix \mathbf{D} , where each edge has integer weight in $[1, U]$, SPARSIFY PRODUCT returns an Eulerian digraph H with $\tilde{O}(m/\varepsilon^2)$ edges such that with high probability*

$$\mathbf{A}_H \stackrel{sv}{\approx}_{\varepsilon} \mathbf{A}_{G_2}\mathbf{D} + \mathbf{A}_{G_1}$$

Moreover, SPARSIFY PRODUCT runs in time $\tilde{O}(m/\varepsilon^2)$.

Algorithm 5: SPARSIFY PRODUCT(ε, G_1, G_2)

1 Let G be the graph with $\mathbf{A}_G = \begin{bmatrix} \mathbf{0} & \mathbf{0} & \mathbf{0} \\ \mathbf{A}_{G_1} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{A}_{G_2} & \mathbf{0} \end{bmatrix}$

2 Let $H \leftarrow [\text{SPARSIFY SQUARE}(\varepsilon/3, G)]_{3,1}$.

3 return H

Proof. We first analyze correctness. Note by Theorem 4.20 we have that

$$\begin{bmatrix} \mathbf{0} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} \\ \mathbf{A}_{H_0} & \mathbf{0} & \mathbf{0} \end{bmatrix} \text{ is an } \varepsilon/6\text{-approximation of } \begin{bmatrix} \mathbf{0} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} \\ \mathbf{A}_{G_2}\mathbf{D} + \mathbf{A}_{G_1} & \mathbf{0} & \mathbf{0} \end{bmatrix} \text{ with respect to } \mathbf{R}$$

where

$$\mathbf{R} = \begin{bmatrix} \mathbf{D} & \mathbf{0} & -(\mathbf{A}_{G_2}\mathbf{D} + \mathbf{A}_{G_1})^\top \\ \mathbf{0} & \mathbf{0} & \mathbf{0} \\ -\mathbf{A}_{G_2}\mathbf{D} + \mathbf{A}_{G_1} & \mathbf{0} & \mathbf{D} \end{bmatrix}$$

This is precisely Item 3 of Lemma 3.7, so we conclude $\mathbf{A}_H \stackrel{sv}{\approx}_{\varepsilon} \mathbf{A}_{G_2}\mathbf{D} + \mathbf{A}_{G_1}$. \square

Finally, we require an algorithm that removes very small edge weights. This is not trivial, as we must do so in a way that exactly preserves degrees for SV approximation to hold.

Lemma 4.22. *There is an algorithm `FIXEDGEWEIGHTS` such that the following holds. Let G be an Eulerian digraph with m edges where the degree matrix \mathbf{D}_G can be represented as a subset sum of $\{1, \dots, 2^b\}$. Then for $t \in \mathbb{N}$, `FIXEDGEWEIGHTS`(t, G) outputs in time $O(mb)$ a graph H such that:*

- H_t has $m + 2n$ edges and is Eulerian and $\mathbf{D}_H = \mathbf{D}_G$.
- Edge weights in H can be represented as a subset sum of $\{2^{-t}, \dots, 1, \dots, 2^b\}$.
- $\|\mathbf{A}_H - \mathbf{A}_G\| \leq 2n \cdot 2^{-t}$.

Proof. First, let $\lfloor G \rfloor$ be the graph where all edge weights are rounded down to a multiple of 2^{-t} . Observe that this satisfies

$$\|\mathbf{A}_{\lfloor G \rfloor} - \mathbf{A}_G\| \leq n \cdot 2^{-t}.$$

Furthermore, note that each vertex v has an in-degree shortfall equal to $s_{in}(v) = \mathbf{D}_{v,v} - \sum_u (\mathbf{A}_{\lfloor G \rfloor})_{v,u}$. Note that $s_{in}(v) \geq 0$ since we only remove weight from edges. This can be expressed as a subset sum of $\{2^{-t}, \dots, 1, \dots, 2^b\}$ as all numbers in the summand can be. An equivalent statement holds for the out-degree shortfall. Furthermore, the total in- and out-degree shortfalls must be equal, since every truncation contributes to both sums. Thus, by greedily matching in- shortfalls to out- shortfalls, we obtain the desired graph H . It is easy to see this satisfies the approximation claim. \square

We are now prepared to state our algorithm for sparsifying an arbitrary power of an Eulerian digraph.

Algorithm 6: `SPARSIFYPOWER`(ε, G, ℓ)

```

1 Let  $t \stackrel{\text{def}}{=} \lceil \log \ell \rceil$ .
2 Let  $\langle \ell \rangle$  be the binary expansion of  $\ell$ .
3 Let  $l \stackrel{\text{def}}{=} O(\log(n\ell/\varepsilon))$ .
4 Let  $G_1 \stackrel{\text{def}}{=} \text{SPARSIFYGRAPH}(\varepsilon/2, G)$ .
5 for  $i = 2, \dots, t$  do
6   | Let  $P_i \stackrel{\text{def}}{=} \text{SPARSIFYPRODUCT}(\varepsilon/t, G_{i-1}, G_{i-1})$ .
7   | Let  $H_i \stackrel{\text{def}}{=} \text{FIXEDGEWEIGHTS}(l, P_i)$ .
8   | Let  $G_i \stackrel{\text{def}}{=} \text{SPARSIFYGRAPH}(\varepsilon/t, H_i)$ .
9   | if  $\langle \ell \rangle_{t-i} = 1$  then
10  |   | Let  $P_i \leftarrow \text{SPARSIFYPRODUCT}(\varepsilon/t, G_{i-1}, G_1)$ .
11  |   | Let  $H_i \leftarrow \text{FIXEDGEWEIGHTS}(l, P_i)$ .
12  |   | Let  $G_i \leftarrow \text{SPARSIFYGRAPH}(\varepsilon/t, H_i)$ .
13  | end
14 end
15 return  $G_t$ 

```

We are now prepared to prove the result:

Theorem 4.17. *Given $\ell \in \mathbb{N}$ and $\varepsilon \in (1/\text{poly}(n), 1)$ and an Eulerian digraph G where $\sigma(G) \leq 1 - 1/\text{poly}(n)$ with m edges with integer edge weights in $[1, \text{poly}(n)]$, SPARSIFYPOWER returns in time $\tilde{O}(m + n\varepsilon^{-2} \log^3(\ell))$ an Eulerian digraph H with $\tilde{O}(n\varepsilon^{-2} \log^2(\ell))$ edges with integer edge weights in $[1, \text{poly}(n, \ell, 1/\varepsilon)]$ such that*

$$\mathbf{D}_H^{+/2} \mathbf{A}_H \mathbf{D}_H^{+/2} \stackrel{svn}{\approx}_{\varepsilon} (\mathbf{D}_G^{+/2} \mathbf{A}_G \mathbf{D}_G^{+/2})^{\ell}.$$

Proof. We first show the approximation guarantee. Note that by the guarantees of Lemma 4.21 and Corollary 4.12 and Lemma 4.22, for every i ,

$$\mathbf{D} \stackrel{\text{def}}{=} \mathbf{D}_{P_i} = \mathbf{D}_{H_i} = \mathbf{D}_{G_i} = \mathbf{D}_G.$$

Let $\mathbf{N}_{P_i} \stackrel{\text{def}}{=} \mathbf{D}^{+/2} \mathbf{A}_{P_i}^{+/2} \mathbf{D}^{+/2}$ and $\mathbf{N}_{H_i} \stackrel{\text{def}}{=} \mathbf{D}^{+/2} \mathbf{A}_{H_i}^{+/2} \mathbf{D}^{+/2}$ and $\mathbf{N}_{G_i} \stackrel{\text{def}}{=} \mathbf{D}^{+/2} \mathbf{A}_{G_i}^{+/2} \mathbf{D}^{+/2}$. The proof of correctness does not change if we assume $\ell = 2^t$ for some t (and hence the if condition in the main loop never occurs), so we analyze it assuming this for simplicity. The fact that this does not matter follows from the fact that SV approximation is preserved under arbitrary products, not merely powers (Lemma 3.16).

We first argue that the singular value of all graphs we sparsify remain bounded below 1.

Claim 4.23. *For every i , $\sigma(H_i), \sigma(P_i), \sigma(G_i) \leq 1 - 1/\text{poly}(n)$.*

Proof. We show the claim via induction. By assumption, we have $\sigma(G) = 1 - 1/n^c$ for some constant c . By Lemma B.4 we have $\sigma(G_1) \leq 1 - (1 + \varepsilon/t)/n^c$. Subsequently, assume that $\sigma(G_{i-1}) \leq 1 - (1 + 2\varepsilon/t)^{2(i-1)}/n^c$.

- By Lemma B.1 we have $\sigma(G_{i-1}^2) \leq \sigma(G_{i-1}) \leq 1 - (1 + 2\varepsilon/t)^{2(i-1)}/n^c$.
- By Lemma B.4 we have $\sigma(P_i) \leq 1 - (1 + \varepsilon/t)(1 + 2\varepsilon/t)^{2(i-1)}/n^c$.
- By Lemma B.3 we have $\sigma(H_i) \leq 1 - (1 + \varepsilon/t)(1 + 2\varepsilon/t)^{2(i-1)}/n^c + 1/n^{10}$.
- By Lemma B.4 we have $\sigma(G_i) \leq 1 - (1 + 2\varepsilon/t)^{2i}/n^c$.

So the claim follows assuming $\varepsilon \leq 1/c'$ for constant c' . □

We now show that SV approximation is preserved at every step of the loop. Fixing i and G_{i-1} :

1. By Lemma 4.21 we have that $\mathbf{N}_{P_i} \stackrel{svn}{\approx}_{\varepsilon/3t} \mathbf{N}_{G_{i-1}}^2$.
2. By Lemma 4.22 we have $\|\mathbf{A}_{H_i} - \mathbf{A}_{P_i}\| \leq 2n2^{-l}$, so \mathbf{A}_{H_i} is a $2n2^{-l} \leq \varepsilon/\text{poly}(tn)$ approximation of \mathbf{A}_{P_i} with respect to \mathbf{D} . Taking the bipartite lift, $\begin{bmatrix} \mathbf{0} & \mathbf{A}_{H_i}^{\top} \\ \mathbf{A}_{H_i} & \mathbf{0} \end{bmatrix}$ is a degree- and bipartition-preserving $\varepsilon/\text{poly}(tn)$ -approximation of $\begin{bmatrix} \mathbf{0} & \mathbf{A}_{P_i}^{\top} \\ \mathbf{A}_{P_i} & \mathbf{0} \end{bmatrix}$ with respect to $\begin{bmatrix} \mathbf{D} & \mathbf{0} \\ \mathbf{0} & \mathbf{D} \end{bmatrix}$ (where we use that P_i is not itself bipartite to ensure that the lift of H_i preserves all bipartitions). Thus by Lemma 4.3 (using that the bipartite lift has normalized second eigenvalue at least $1/\text{poly}(n)$ by Claim 4.23) and Lemma 3.7, we have that $\mathbf{N}_{H_i} \stackrel{sv}{\approx}_{\varepsilon/3t} \mathbf{N}_{P_i}$.
3. By Corollary 4.12, $\mathbf{N}_{G_i} \stackrel{svn}{\approx}_{\varepsilon/3t} \mathbf{N}_{H_i}$.

Applying Lemma 3.19 we obtain $\mathbf{N}_{G_i} \stackrel{svn}{\approx}_{\varepsilon/t} \mathbf{N}_{G_{i-1}}^2$ for every i . Therefore by Lemma 3.16,

$$\mathbf{N}_{G_t} \stackrel{svn}{\approx}_{\varepsilon/t} (\mathbf{N}_{G_{i-1}})^2 \stackrel{svn}{\approx}_{\varepsilon/t} \dots (\mathbf{N}_{G_2})^{2^{t-1}} \stackrel{svn}{\approx}_{\varepsilon/t} (\mathbf{N}_{G_1})^{2^t}$$

and so we conclude by Lemma 3.19 (and taking $\varepsilon \leftarrow \varepsilon/10$).

To analyze the time, note that by Lemma 4.22 we always call SPARSIFYGRAPH on graphs with (up to rescaling) integer edge weights in $[1, \text{poly}(n\ell/\varepsilon)]$, and by Theorem 4.20 P_i (and thus H_i) has $\tilde{O}(n\varepsilon^{-2} \log^2 \ell)$ edges, so we obtain the desired runtime and sparsity by Corollary 4.12. \square

5 Squaring algorithm for solving normal directed Laplacians

In [PS14], Peng and Spielman proposed a squaring-based parallel algorithm for computing an approximate pseudo-inverse (preconditioner) of a Laplacian matrix. A key ingredient of their algorithm is the following recursion, which we refer to as the *Peng-Spielman squaring recursion* (*PS-recursion*), for inverting $\mathbf{I} - \mathbf{W}$ when $\|\mathbf{W}\| < 1$.⁶

$$(\mathbf{I} - \mathbf{W})^{-1} = \frac{1}{2} [\mathbf{I} + (\mathbf{I} + \mathbf{W})(\mathbf{I} - \mathbf{W}^2)^{-1}(\mathbf{I} + \mathbf{W})] \quad (19)$$

Leveraging the PS-recursion a natural approach to find an approximate inverse of $\mathbf{I} - \mathbf{W}$, is to compute a sparse matrix $\widetilde{\mathbf{W}}$ such that $\mathbf{I} - \widetilde{\mathbf{W}} \approx \mathbf{I} - \mathbf{W}^2$, and then compute the inverse of $\mathbf{I} - \widetilde{\mathbf{W}}$ recursively. [PS14] showed that for symmetric Laplacians, using spectral approximation this recursion can be leveraged to produce a constant preconditioner with logarithmic depth.

Here we prove that the same algorithm gives approximate pre-conditioner for $\mathbf{I} - \mathbf{W}$ if \mathbf{W} is normal but not necessarily symmetric and the square approximations are normalized SV approximations. We think Theorem 5.2 gives hope that a similar approach can expand the proof to non-normal matrices. If proved, this will result in simple but space and time efficient algorithms for solving directed Laplacian systems. For example, this could simplify the space efficient solver introduced in [AKM⁺20] which in turn is used to estimate random walk probabilities to a high precision.

Below we give the main theorem which formally defines the PS-recursion and states the main result of this section. This theorem uses a generalization of the spectral norm to a (semi-)norm induced by a PSD matrix, which we first define.

Definition 5.1. Given PSD $\mathbf{F} \in \mathbb{C}^{n \times n}$, we let $\|\cdot\|_{\mathbf{F}}$ be the (semi)norm on vectors where $\|x\|_{\mathbf{F}} \stackrel{\text{def}}{=} \sqrt{x^* \mathbf{F} x}$, and define the matrix norm $\|\mathbf{A}\|_{\mathbf{F}} \stackrel{\text{def}}{=} \sup_{x \in \mathbb{C}^n \setminus \{\vec{0}\}} \frac{\|\mathbf{A}x\|_{\mathbf{F}}}{\|x\|_{\mathbf{F}}}$.

Theorem 5.2. For a normal matrix $\mathbf{W} \in \mathbb{C}^{n \times n}$ with $\|\mathbf{W}\| \leq 1$, let $\mathbf{W} = \mathbf{W}_0, \dots, \mathbf{W}_{k-1}$ be a sequence of matrices such that for all $i \in [k-1]$ we have $\mathbf{W}_i \stackrel{svn}{\approx}_{\varepsilon} \mathbf{W}_{i-1}^2$ for $\varepsilon \leq 1/(4k)$. Define

$$\mathbf{P}_i \stackrel{\text{def}}{=} \frac{1}{2} [\mathbf{I} + (\mathbf{I} + \mathbf{W}_i)\mathbf{P}_{i+1}(\mathbf{I} + \mathbf{W}_i)] \quad \text{for all } 0 \leq i < k$$

where \mathbf{P}_k is a matrix such that

$$\left\| (\mathbf{I} - \mathbf{W}^{2^k})^{\frac{1}{2}} \left[\mathbf{P}_k - (\mathbf{I} - \mathbf{W}^{2^k})^+ \right] (\mathbf{I} - \mathbf{W}^{2^k})^{\frac{1}{2}} \right\| \leq O(k\varepsilon).$$

Then for $\mathbf{B} \stackrel{\text{def}}{=} ((\mathbf{I} - \mathbf{W})^{1/2})^* (\mathbf{I} - \mathbf{W})^{1/2}$

$$\|\mathbf{P}_0(\mathbf{I} - \mathbf{W}) - \mathbf{I}\|_{\mathbf{B}} = \left\| (\mathbf{I} - \mathbf{W})^{\frac{1}{2}} \left[\mathbf{P}_0 - (\mathbf{I} - \mathbf{W})^+ \right] (\mathbf{I} - \mathbf{W})^{\frac{1}{2}} \right\| \leq O(k^2\varepsilon). \quad (20)$$

⁶Technically the recursion was introduced for symmetric \mathbf{W} .

Note that in the above statement, the square root of $\mathbf{I} - \mathbf{W}$ is well defined by the assumption that \mathbf{W} is normal. In the case of non-normal matrices, which is outside the scope of this paper, one can consider the Maclaurin series for $(1 - z)^{1/2}$ to work with $(\mathbf{I} - \mathbf{W})^{1/2}$.

We do not give a full algorithm for solving systems defined by Theorem 5.2 as the main purpose is to show that the error analysis of the squaring algorithm can be expanded to normal asymmetric matrices, and we already know space and time efficient algorithms [CKK⁺18, AKM⁺20] for solving a larger class of matrices which includes non-normal \mathbf{W} 's. However, one can still get an actual squaring based solver from Theorem 5.2. For example, for a normal \mathbf{W} corresponding to a random walk matrix of an Eulerian digraph one can carry the same steps in [AKM⁺20] by setting $k = O(\log n)$ and $\mathbf{P}_k = \mathbf{I} - \frac{\overline{\mathbf{1}\mathbf{1}^\top}}{n}$ to get an actual squaring based solver using Theorem 5.2 and the derandomized square sparsification described in Section 4.5.

We prove the statement of the theorem, following a few helper lemmas. First, we show the error of \mathbf{W}_i 's approximating \mathbf{W}^{2^i} 's grows additively.

Lemma 5.3. *Let \mathbf{W}_i be defined as in Theorem 5.2. Then $\mathbf{W}_i \stackrel{svn}{\approx}_{2i\epsilon} \mathbf{W}^{2^i}$ for all for $i \in [k]$.*

Proof. Recall that $\mathbf{W}_j \stackrel{svn}{\approx}_\epsilon \mathbf{W}_{j-1}^2$ for every j . Thus, by Lemma 3.16 we have for every $j \leq i$,

$$(\mathbf{W}_j)^{2^{i-j}} \stackrel{svn}{\approx}_{\epsilon + O(\epsilon^2)} (\mathbf{W}_{j-1})^{2^{i-j+1}}.$$

The result then follows from Lemma 3.19. \square

Next we give a general technical tool regarding normal matrices.

Lemma 5.4. *For a normal matrix \mathbf{V} with $\|\mathbf{V}\| \leq 1$*

$$\|(\mathbf{I} + \mathbf{V})^{\frac{1}{2}}\| \leq \sqrt{2} \tag{21}$$

and if $\tilde{\mathbf{V}} \stackrel{svn}{\approx}_\delta \mathbf{V}$ then

$$\left\| (\mathbf{I} - \mathbf{V})^{\frac{1}{2}} (\tilde{\mathbf{V}} - \mathbf{V}) ((\mathbf{I} - \mathbf{V}^2)^+)^{\frac{1}{2}} \right\| \leq O(\delta). \tag{22}$$

Proof. Note that since \mathbf{V} is normal, we can write it as $\mathbf{V} = \mathbf{U}\mathbf{D}\mathbf{U}^*$ where \mathbf{U} is unitary and \mathbf{D} is diagonal and $\|\mathbf{D}\| \leq 1$. Then it is easy to see

$$\|(\mathbf{I} + \mathbf{V})^{\frac{1}{2}}\| = \|\mathbf{U}(\mathbf{I} + \mathbf{D})^{\frac{1}{2}}\mathbf{U}^*\| = \|(\mathbf{I} + \mathbf{D})^{\frac{1}{2}}\| \leq \sqrt{2}. \tag{23}$$

To prove (22), given the SV approximation definition, for unit vectors x, y we have,

$$\begin{aligned} \left| x^* (\mathbf{I} - \mathbf{V})^{\frac{1}{2}} (\tilde{\mathbf{V}} - \mathbf{V}) ((\mathbf{I} - \mathbf{V}^2)^+)^{\frac{1}{2}} y \right| &\leq \frac{\delta}{2} \cdot x^* (\mathbf{I} - \mathbf{V})^{\frac{1}{2}} (\mathbf{I} - \mathbf{V}\mathbf{V}^*) (\mathbf{I} - \mathbf{V}^*)^{\frac{1}{2}} x \\ &\quad + \frac{\delta}{2} \cdot y^* ((\mathbf{I} - (\mathbf{V}^*)^2)^+)^{\frac{1}{2}} (\mathbf{I} - \mathbf{V}^*\mathbf{V}) ((\mathbf{I} - \mathbf{V}^2)^+)^{\frac{1}{2}} y \end{aligned}$$

We bound each term on the right hand side separately. Since \mathbf{V} is normal we can prove bounds on the norm of $(\mathbf{I} - \mathbf{V})^{\frac{1}{2}} (\mathbf{I} - \mathbf{V}\mathbf{V}^*) (\mathbf{I} - \mathbf{V}^*)^{\frac{1}{2}}$ and $((\mathbf{I} - (\mathbf{V}^*)^2)^+)^{\frac{1}{2}} (\mathbf{I} - \mathbf{V}^*\mathbf{V}) ((\mathbf{I} - \mathbf{V}^2)^+)^{\frac{1}{2}}$ by looking at complex numbers. Note that for $z \in \mathbb{C}$ and $|z| \leq 1$,

$$\left| (1 - z)^{\frac{1}{2}} (1 - z \cdot z^*)^2 (1 - z^*)^{\frac{1}{2}} \right| = |1 - z| \cdot (1 - |z|^2) \leq 2,$$

thus $\|(\mathbf{I} - \mathbf{V})^{\frac{1}{2}} (\mathbf{I} - \mathbf{V}\mathbf{V}^*) (\mathbf{I} - \mathbf{V}^*)^{\frac{1}{2}}\| \leq 2$. Next,

$$\left| (1 - z^{*2})^{-\frac{1}{2}} (1 - z \cdot z^*)^2 (1 - z^2)^{-\frac{1}{2}} \right| = \frac{1 - |z|^2}{|1 - z^2|} \leq 1$$

because $1 - |z|^2 \leq |1 - z^2|$ by triangle inequality. Thus,

$$\|((\mathbf{I} - (\mathbf{V}^*)^2)^+)^{\frac{1}{2}}(\mathbf{I} - \mathbf{V}^*\mathbf{V})((\mathbf{I} - \mathbf{V}^2)^+)^{\frac{1}{2}}\| \leq 1.$$

Completing the proof that,

$$\left| x^*(\mathbf{I} - \mathbf{V})^{\frac{1}{2}}(\tilde{\mathbf{V}} - \mathbf{V})((\mathbf{I} - \mathbf{V}^2)^+)^{\frac{1}{2}}y \right| \leq \frac{3}{2} \cdot \delta.$$

□

Leveraging these tools we give the central lemma towards proving Theorem 5.2 and conclude by proving Theorem 5.2.

Lemma 5.5. *For ϵ and \mathbf{P}_i 's defined as in Theorem 5.2,*

$$\left\| (\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}} \left[\mathbf{P}_i - (\mathbf{I} - \mathbf{W}^{2^i})^+ \right] (\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}} \right\| \leq \epsilon_i \quad (24)$$

where $\epsilon_i = (1 + O(i \cdot \epsilon))\epsilon_{i+1} + O(i \cdot \epsilon)$ for $0 \leq i < k$ and $\epsilon_k = O(k \cdot \epsilon)$.

Proof. We prove the statement by backward induction. Note that the statement is true for $i = k$ by the assumption about \mathbf{P}_k in Theorem 5.2. Let

$$\mathbf{Q}_i = \frac{1}{2} \left[\mathbf{I} + (\mathbf{I} + \mathbf{W}_i)(\mathbf{I} - \mathbf{W}^{2^{i+1}})^+(\mathbf{I} + \mathbf{W}_i) \right],$$

$$\mathbf{R}_i = \frac{1}{2} \left[\mathbf{I} + (\mathbf{I} + \mathbf{W}_i)(\mathbf{I} - \mathbf{W}^{2^{i+1}})^+(\mathbf{I} + \mathbf{W}^{2^i}) \right],$$

$$\mathbf{S}_i = \frac{1}{2} \left[\mathbf{I} + (\mathbf{I} + \mathbf{W}^{2^i})(\mathbf{I} - \mathbf{W}^{2^{i+1}})^+(\mathbf{I} + \mathbf{W}^{2^i}) \right] = (\mathbf{I} - \mathbf{W}^{2^i})^+.$$

Then by triangle inequality we have,

$$\|(\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}} \left[\mathbf{P}_i - (\mathbf{I} - \mathbf{W}^{2^i})^+ \right] (\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}}\| \leq \|(\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}}(\mathbf{P}_i - \mathbf{Q}_i)(\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}}\| \quad (25)$$

$$+ \|(\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}}(\mathbf{Q}_i - \mathbf{R}_i)(\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}}\| \quad (26)$$

$$+ \|(\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}}(\mathbf{R}_i - \mathbf{S}_i)(\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}}\|. \quad (27)$$

Next, we bound each term separately. For (25), we have

$$\begin{aligned} & \left\| (\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}}(\mathbf{P}_i - \mathbf{Q}_i)(\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}} \right\| \\ &= \frac{1}{2} \left\| (\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}}(\mathbf{I} + \mathbf{W}_i) \left(\mathbf{P}_{i+1} - (\mathbf{I} - \mathbf{W}^{2^{i+1}})^+ \right) (\mathbf{I} + \mathbf{W}_i)(\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}} \right\| \\ &\leq \frac{1}{2} \left\| (\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}}(\mathbf{I} + \mathbf{W}_i)((\mathbf{I} - \mathbf{W}^{2^{i+1}})^+)^{\frac{1}{2}} \right\| \\ &\quad \cdot \left\| (\mathbf{I} - \mathbf{W}^{2^{i+1}})^{\frac{1}{2}} \left(\mathbf{P}_{i+1} - (\mathbf{I} - \mathbf{W}^{2^{i+1}})^+ \right) (\mathbf{I} - \mathbf{W}^{2^{i+1}})^{\frac{1}{2}} \right\| \\ &\quad \cdot \left\| ((\mathbf{I} - \mathbf{W}^{2^{i+1}})^+)^{\frac{1}{2}}(\mathbf{I} + \mathbf{W}_i)(\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}} \right\|. \end{aligned}$$

Note that the middle term above satisfies

$$\left\| (\mathbf{I} - \mathbf{W}^{2^{i+1}})^{\frac{1}{2}} \left(\mathbf{P}_{i+1} - (\mathbf{I} - \mathbf{W}^{2^{i+1}})^+ \right) (\mathbf{I} - \mathbf{W}^{2^{i+1}})^{\frac{1}{2}} \right\| \leq \epsilon_{i+1}$$

by the induction hypothesis. Next, we bound the first and third terms similarly:

$$\begin{aligned} \left\| \frac{(\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}}(\mathbf{I} + \mathbf{W}_i)((\mathbf{I} - \mathbf{W}^{2^{i+1}})^{\frac{1}{2}})^+}{\sqrt{2}} \right\| &\leq \left\| \frac{(\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}}(\mathbf{I} + \mathbf{W}^{2^i})((\mathbf{I} - \mathbf{W}^{2^{i+1}})^{\frac{1}{2}})^+}{\sqrt{2}} \right\| + \\ &\left\| \frac{(\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}}(\mathbf{W}_i - \mathbf{W}^{2^i})((\mathbf{I} - \mathbf{W}^{2^{i+1}})^{\frac{1}{2}})^+}{\sqrt{2}} \right\| \\ &\leq 1 + O(i \cdot \epsilon) \end{aligned} \quad (28)$$

as $\left\| \frac{(\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}}(\mathbf{I} + \mathbf{W}^{2^i})((\mathbf{I} - \mathbf{W}^{2^{i+1}})^{\frac{1}{2}})^+}{\sqrt{2}} \right\| \leq 1$ by Equation (21) in Lemma 5.4 and $\left\| \frac{(\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}}(\mathbf{W}_i - \mathbf{W}^{2^i})((\mathbf{I} - \mathbf{W}^{2^{i+1}})^{\frac{1}{2}})^+}{\sqrt{2}} \right\| \leq O(i \cdot \epsilon)$ by Lemma 5.3 and Equation (22) in Lemma 5.4. Combining all the above,

$$\left\| (\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}}(\mathbf{P}_i - \mathbf{Q}_i)(\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}} \right\| \leq (1 + O(i \cdot \epsilon))^2 \epsilon_{i+1},$$

giving the bound for (25). Next, for (26), we have

$$\begin{aligned} &\left\| (\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}}(\mathbf{Q}_i - \mathbf{R}_i)(\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}} \right\| = \\ &\frac{1}{2} \left\| (\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}}(\mathbf{I} + \mathbf{W}_i)(\mathbf{I} - \mathbf{W}^{2^{i+1}})^+ (\mathbf{W}_i - \mathbf{W}^{2^i})(\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}} \right\| \leq \\ &\left\| \frac{(\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}}(\mathbf{I} + \mathbf{W}_i)((\mathbf{I} - \mathbf{W}^{2^{i+1}})^{\frac{1}{2}})^+}{\sqrt{2}} \right\| \cdot \left\| \frac{((\mathbf{I} - \mathbf{W}^{2^{i+1}})^{\frac{1}{2}})^+ (\mathbf{W}_i - \mathbf{W}^{2^i})(\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}}}{\sqrt{2}} \right\| \end{aligned}$$

We can get bounds for the two terms above similar to (28). Thus we get,

$$\begin{aligned} &\left\| (\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}}(\mathbf{Q}_i - \mathbf{R}_i)(\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}} \right\| \leq \\ &\left\| \frac{(\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}}(\mathbf{I} + \mathbf{W}_i)((\mathbf{I} - \mathbf{W}^{2^{i+1}})^{\frac{1}{2}})^+}{\sqrt{2}} \right\| \cdot \left\| \frac{((\mathbf{I} - \mathbf{W}^{2^{i+1}})^{\frac{1}{2}})^+ (\mathbf{W}_i - \mathbf{W}^{2^i})(\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}}}{\sqrt{2}} \right\| \leq \\ &(1 + O(i \cdot \epsilon)) \cdot O(i \cdot \epsilon) \leq O(i \cdot \epsilon). \end{aligned}$$

Finally, for (27), we have

$$\begin{aligned} &\left\| (\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}}(\mathbf{R}_i - \mathbf{S}_i)(\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}} \right\| = \\ &\frac{1}{2} \left\| (\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}}(\mathbf{W}_i - \mathbf{W}^{2^i})(\mathbf{I} - \mathbf{W}^{2^{i+1}})^+ (\mathbf{I} + \mathbf{W}^{2^i})(\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}} \right\| \leq \\ &\left\| \frac{(\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}}(\mathbf{W}_i - \mathbf{W}^{2^i})((\mathbf{I} - \mathbf{W}^{2^{i+1}})^{\frac{1}{2}})^+}{\sqrt{2}} \right\| \cdot \left\| \frac{((\mathbf{I} - \mathbf{W}^{2^{i+1}})^{\frac{1}{2}})^+ (\mathbf{I} + \mathbf{W}^{2^i})(\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}}}{\sqrt{2}} \right\| \end{aligned}$$

Further, similar to the earlier upper bounds, we have $\left\| \frac{(\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}}(\mathbf{W}_i - \mathbf{W}^{2^i})((\mathbf{I} - \mathbf{W}^{2^{i+1}})^{\frac{1}{2}})^+}{\sqrt{2}} \right\| \leq O(i \cdot \epsilon)$

and $\left\| \frac{((\mathbf{I} - \mathbf{W}^{2^{i+1}})^{\frac{1}{2}})^+ (\mathbf{I} + \mathbf{W}^{2^i})(\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}}}{\sqrt{2}} \right\| \leq 1$. Thus

$$\left\| (\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}}(\mathbf{R}_i - \mathbf{S}_i)(\mathbf{I} - \mathbf{W}^{2^i})^{\frac{1}{2}} \right\| \leq O(i \cdot \epsilon).$$

□

Proof of Theorem 5.2. For $\epsilon < \frac{1}{k^2}$, we can solve for ϵ_0 from $\epsilon_i = (1 + O(i \cdot \epsilon))\epsilon_{i+1} + O(i \cdot \epsilon)$ in Lemma 5.5 and get $\epsilon_0 = O(\epsilon \cdot k^2)$. To complete the proof, we show

$$\|\mathbf{P}_0(\mathbf{I} - \mathbf{W}) - \mathbf{I}\|_{\mathbf{B}} = \|(\mathbf{I} - \mathbf{W})^{\frac{1}{2}} [\mathbf{P}_0 - (\mathbf{I} - \mathbf{W})^+] (\mathbf{I} - \mathbf{W})^{\frac{1}{2}}\|.$$

Note that,

$$\begin{aligned} & \|(\mathbf{I} - \mathbf{W})^{\frac{1}{2}} [\mathbf{P}_0 - (\mathbf{I} - \mathbf{W})^+] (\mathbf{I} - \mathbf{W})^{\frac{1}{2}}\| \\ &= \|(\mathbf{I} - \mathbf{W})^{1/2} [\mathbf{P}_0(\mathbf{I} - \mathbf{W}) - \mathbf{I}] (\mathbf{I} - \mathbf{W})^{+2}\| \\ &= \left\| \left[(\mathbf{I} - \mathbf{W})^{*/2} (\mathbf{I} - \mathbf{W})^{1/2} \right]^{1/2} [\mathbf{P}_0(\mathbf{I} - \mathbf{W}) - \mathbf{I}] \left[(\mathbf{I} - \mathbf{W})^{+2} ((\mathbf{I} - \mathbf{W})^*)^{+2} \right]^{1/2} \right\| \\ &= \left\| \left[(\mathbf{I} - \mathbf{W})^{*/2} (\mathbf{I} - \mathbf{W})^{1/2} \right]^{1/2} [\mathbf{P}_0(\mathbf{I} - \mathbf{W}) - \mathbf{I}] \left[(\mathbf{I} - \mathbf{W})^{*/2} (\mathbf{I} - \mathbf{W})^{1/2} \right]^{+2} \right\| \\ &= \|\mathbf{P}_0(\mathbf{I} - \mathbf{W}) - \mathbf{I}\|_{\mathbf{B}} \end{aligned}$$

where we used Lemma B.6. □

Acknowledgements

We thank Jack Murtagh for his helpful collaboration at the early stage of research on this research and we thank Jonathan Kelner for helpful conversations at various stages of work on this project.

Aaron Sidford was supported by a Microsoft Research Faculty Fellowship, NSF CAREER Award CCF-1844855, NSF Grant CCF-1955039, a PayPal research award, and a Sloan Research Fellowship.

References

- [ABKS21] Deeksha Adil, Brian Bullins, Rasmus Kyng, and Sushant Sachdeva. Almost-linear-time weighted lp-norm solvers in slightly dense graphs via sparsification. In Nikhil Bansal, Emanuela Merelli, and James Worrell, editors, *48th International Colloquium on Automata, Languages, and Programming, ICALP 2021, July 12-16, 2021, Glasgow, Scotland (Virtual Conference)*, volume 198 of *LIPICs*, pages 9:1–9:15. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2021.
- [ACL06] Reid Andersen, Fan R. K. Chung, and Kevin J. Lang. Local graph partitioning using pagerank vectors. In *47th Annual IEEE Symposium on Foundations of Computer Science (FOCS 2006), 21-24 October 2006, Berkeley, California, USA, Proceedings*, pages 475–486. IEEE Computer Society, 2006.
- [ADK22] Daniel Agassy, Dani Dorfman, and Haim Kaplan. Expander decomposition with fewer inter-cluster edges using a spectral cut player. *CoRR*, abs/2205.10301, 2022.
- [AG15] Nima Anari and Shayan Oveis Gharan. Effective-resistance-reducing flows, spectrally thin trees, and asymmetric TSP. In Venkatesan Guruswami, editor, *IEEE 56th Annual Symposium on Foundations of Computer Science, FOCS 2015, Berkeley, CA, USA, 17-20 October, 2015*, pages 20–39. IEEE Computer Society, 2015.

- [AGM⁺10] Arash Asadpour, Michel X. Goemans, Aleksander Madry, Shayan Oveis Gharan, and Amin Saberi. An $o(\log n / \log \log n)$ -approximation algorithm for the asymmetric traveling salesman problem. In Moses Charikar, editor, *Proceedings of the Twenty-First Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2010, Austin, Texas, USA, January 17-19, 2010*, pages 379–389. SIAM, 2010.
- [Ahm20] AmirMahdi Ahmadinejad. *Computing Stationary Distributions: Perron Vectors, Random Walks, and Ride-Sharing Competition*. Stanford University, 2020.
- [AJSS19] AmirMahdi Ahmadinejad, Arun Jambulapati, Amin Saberi, and Aaron Sidford. Perron-frobenius theory in nearly linear time: Positive eigenvectors, m-matrices, graph kernels, and other applications. In *Proceedings of the Thirtieth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2019, San Diego, California, USA, January 6-9, 2019*, pages 1387–1404. SIAM, 2019.
- [AKM⁺20] AmirMahdi Ahmadinejad, Jonathan A. Kelner, Jack Murtagh, John Peebles, Aaron Sidford, and Salil P. Vadhan. High-precision estimation of random walks in small space. In Sandy Irani, editor, *61st IEEE Annual Symposium on Foundations of Computer Science, FOCS 2020, Durham, NC, USA, November 16-19, 2020*, pages 1295–1306. IEEE, 2020.
- [AM85] Noga Alon and V. D. Milman. λ_1 , isoperimetric inequalities for graphs, and superconcentrators. *J. Comb. Theory, Ser. B*, 38(1):73–88, 1985.
- [BK00] András A. Benczúr and David R. Karger. Augmenting undirected edge connectivity in $\tilde{O}(n^2)$ time. *J. Algorithms*, 37(1):2–36, 2000.
- [BSS12] Joshua D. Batson, Daniel A. Spielman, and Nikhil Srivastava. Twice-ramanujan sparsifiers. *SIAM J. Comput.*, 41(6):1704–1721, 2012.
- [CGP⁺18] Timothy Chu, Yu Gao, Richard Peng, Sushant Sachdeva, Saurabh Sawlani, and Junxing Wang. Graph sparsification, spectral sketches, and faster resistance computation, via short cycle decompositions. In Mikkel Thorup, editor, *59th IEEE Annual Symposium on Foundations of Computer Science, FOCS 2018, Paris, France, October 7-9, 2018*, pages 361–372. IEEE Computer Society, 2018.
- [CKK⁺18] Michael B. Cohen, Jonathan A. Kelner, Rasmus Kyng, John Peebles, Richard Peng, Anup B. Rao, and Aaron Sidford. Solving directed laplacian systems in nearly-linear time through sparse LU factorizations. In Mikkel Thorup, editor, *59th IEEE Annual Symposium on Foundations of Computer Science, FOCS 2018, Paris, France, October 7-9, 2018*, pages 898–909. IEEE Computer Society, 2018.
- [CKM⁺11] Paul F. Christiano, Jonathan A. Kelner, Aleksander Madry, Daniel A. Spielman, and Shang-Hua Teng. Electrical flows, laplacian systems, and faster approximation of maximum flow in undirected graphs. In Lance Fortnow and Salil P. Vadhan, editors, *Proceedings of the 43rd ACM Symposium on Theory of Computing, STOC 2011, San Jose, CA, USA, 6-8 June 2011*, pages 273–282. ACM, 2011.
- [CKM⁺14] Michael B. Cohen, Rasmus Kyng, Gary L. Miller, Jakub W. Pachocki, Richard Peng, Anup B. Rao, and Shen Chen Xu. Solving SDD linear systems in nearly $m \log^{1/2} n$ time. In David B. Shmoys, editor, *Symposium on Theory of Computing, STOC 2014, New York, NY, USA, May 31 - June 03, 2014*, pages 343–352. ACM, 2014.

- [CKP⁺16] Michael B. Cohen, Jonathan A. Kelner, John Peebles, Richard Peng, Aaron Sidford, and Adrian Vladu. Faster algorithms for computing the stationary distribution, simulating random walks, and more. In Irit Dinur, editor, *IEEE 57th Annual Symposium on Foundations of Computer Science, FOCS 2016, 9-11 October 2016, Hyatt Regency, New Brunswick, New Jersey, USA*, pages 583–592. IEEE Computer Society, 2016.
- [CKP⁺17] Michael B. Cohen, Jonathan A. Kelner, John Peebles, Richard Peng, Anup B. Rao, Aaron Sidford, and Adrian Vladu. Almost-linear-time algorithms for markov chains and new spectral primitives for directed graphs. In Hamed Hatami, Pierre McKenzie, and Valerie King, editors, *Proceedings of the 49th Annual ACM SIGACT Symposium on Theory of Computing, STOC 2017, Montreal, QC, Canada, June 19-23, 2017*, pages 410–419. ACM, 2017.
- [CLTW22] Lijie Chen, Xin Lyu, Avishay Tal, and Hongxun Wu. Personal communication, November 2022.
- [GV22] Louis Golowich and Salil Vadhan. Pseudorandomness of expander random walks for symmetric functions and permutation branching programs. In Anindya De, editor, *Proceedings of the 37th Computational Complexity Conference (CCC ‘22)*, volume 234 of *LIPICs*, pages 27:1–27:15. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2022. Full version posted as ECCO TR22-024.
- [HPV21] William M. Hoza, Edward Pyne, and Salil P. Vadhan. Pseudorandom generators for unbounded-width permutation branching programs. In James R. Lee, editor, *12th Innovations in Theoretical Computer Science Conference, ITCS 2021, January 6-8, 2021, Virtual Conference*, volume 185 of *LIPICs*, pages 7:1–7:20. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2021.
- [INW94] Russell Impagliazzo, Noam Nisan, and Avi Wigderson. Pseudorandomness for network algorithms. In Frank Thomson Leighton and Michael T. Goodrich, editors, *Proceedings of the Twenty-Sixth Annual ACM Symposium on Theory of Computing, 23-25 May 1994, Montréal, Québec, Canada*, pages 356–364. ACM, 1994.
- [JS21] Arun Jambulapati and Aaron Sidford. Ultrasparse ultrasparsifiers and faster laplacian system solvers. In Dániel Marx, editor, *Proceedings of the 2021 ACM-SIAM Symposium on Discrete Algorithms, SODA 2021, Virtual Conference, January 10 - 13, 2021*, pages 540–559. SIAM, 2021.
- [Kel21] Zander Kelley. An alternative proof that inw fools permutation programs. Unpublished manuscript, February 2021.
- [KLOS14] Jonathan A. Kelner, Yin Tat Lee, Lorenzo Orecchia, and Aaron Sidford. An almost-linear-time algorithm for approximate max flow in undirected graphs, and its multi-commodity generalizations. In Chandra Chekuri, editor, *Proceedings of the Twenty-Fifth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2014, Portland, Oregon, USA, January 5-7, 2014*, pages 217–226. SIAM, 2014.
- [KM09] Jonathan A. Kelner and Aleksander Madry. Faster generation of random spanning trees. In *50th Annual IEEE Symposium on Foundations of Computer Science, FOCS 2009, October 25-27, 2009, Atlanta, Georgia, USA*, pages 13–21. IEEE Computer Society, 2009.

- [KMG22] Rasmus Kyng, Simon Meierhans, and Maximilian Probst Gutenberg. Derandomizing directed random walks in almost-linear time. *CoRR*, abs/2208.10959, 2022.
- [KMP12] Jonathan A. Kelner, Gary L. Miller, and Richard Peng. Faster approximate multi-commodity flow using quadratically coupled flows. In Howard J. Karloff and Toniann Pitassi, editors, *Proceedings of the 44th Symposium on Theory of Computing Conference, STOC 2012, New York, NY, USA, May 19 - 22, 2012*, pages 1–18. ACM, 2012.
- [KMP14] Ioannis Koutis, Gary L. Miller, and Richard Peng. Approaching optimality for solving SDD linear systems. *SIAM J. Comput.*, 43(1):337–354, 2014.
- [KVV04] Ravi Kannan, Santosh S. Vempala, and Adrian Vetta. On clusterings: Good, bad and spectral. *J. ACM*, 51(3):497–515, 2004.
- [LRS13] Yin Tat Lee, Satish Rao, and Nikhil Srivastava. A new approach to computing maximum flows using electrical flows. In Dan Boneh, Tim Roughgarden, and Joan Feigenbaum, editors, *Symposium on Theory of Computing Conference, STOC’13, Palo Alto, CA, USA, June 1-4, 2013*, pages 755–764. ACM, 2013.
- [LS13] Yin Tat Lee and Aaron Sidford. Efficient accelerated coordinate descent methods and faster algorithms for solving linear systems. In *54th Annual IEEE Symposium on Foundations of Computer Science, FOCS 2013, 26-29 October, 2013, Berkeley, CA, USA*, pages 147–156. IEEE Computer Society, 2013.
- [LS18] Yin Tat Lee and He Sun. Constructing linear-sized spectral sparsification in almost-linear time. *SIAM J. Comput.*, 47(6):2315–2336, 2018.
- [MST15] Aleksander Madry, Damian Straszak, and Jakub Tarnawski. Fast generation of random spanning trees and the effective resistance metric. In Piotr Indyk, editor, *Proceedings of the Twenty-Sixth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2015, San Diego, CA, USA, January 4-6, 2015*, pages 2019–2036. SIAM, 2015.
- [OSV12] Lorenzo Orecchia, Sushant Sachdeva, and Nisheeth K. Vishnoi. Approximating the exponential, the lanczos method and an $\tilde{o}(m)$ -time spectral algorithm for balanced separator. In Howard J. Karloff and Toniann Pitassi, editors, *Proceedings of the 44th Symposium on Theory of Computing Conference, STOC 2012, New York, NY, USA, May 19 - 22, 2012*, pages 1141–1160. ACM, 2012.
- [PS14] Richard Peng and Daniel A. Spielman. An efficient parallel solver for SDD linear systems. *STOC*, 2014.
- [PV21] Edward Pyne and Salil Vadhan. Pseudodistributions That Beat All Pseudorandom Generators (Extended Abstract). In *Proceedings of the 36th Annual Computational Complexity Conference (CCC)*, pages 33:1–33:15, 2021.
- [PY19] Merav Parter and Eylon Yogev. Optimal short cycle decomposition in almost linear time. In Christel Baier, Ioannis Chatzigiannakis, Paola Flocchini, and Stefano Leonardi, editors, *46th International Colloquium on Automata, Languages, and Programming, ICALP 2019, July 9-12, 2019, Patras, Greece*, volume 132 of *LIPIcs*, pages 89:1–89:14. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2019.

- [RV05] Eyal Rozenman and Salil P. Vadhan. Derandomized squaring of graphs. In Chandra Chekuri, Klaus Jansen, José D. P. Rolim, and Luca Trevisan, editors, *Approximation, Randomization and Combinatorial Optimization, Algorithms and Techniques, 8th International Workshop on Approximation Algorithms for Combinatorial Optimization Problems, APPROX 2005 and 9th International Workshop on Randomization and Computation, RANDOM 2005, Berkeley, CA, USA, August 22-24, 2005, Proceedings*, volume 3624 of *Lecture Notes in Computer Science*, pages 436–447. Springer, 2005.
- [SS08] Daniel A. Spielman and Nikhil Srivastava. Graph sparsification by effective resistances. In Cynthia Dwork, editor, *Proceedings of the 40th Annual ACM Symposium on Theory of Computing, Victoria, British Columbia, Canada, May 17-20, 2008*, pages 563–568. ACM, 2008.
- [ST04] Daniel A. Spielman and Shang-Hua Teng. Nearly-linear time algorithms for graph partitioning, graph sparsification, and solving linear systems. In László Babai, editor, *Proceedings of the 36th Annual ACM Symposium on Theory of Computing, Chicago, IL, USA, June 13-16, 2004*, pages 81–90. ACM, 2004.
- [Tro12] Joel A. Tropp. User-friendly tail bounds for sums of random matrices. *Found. Comput. Math.*, 12(4):389–434, 2012.
- [Vad12] Salil P. Vadhan. Pseudorandomness. *Found. Trends Theor. Comput. Sci.*, 7(1-3):1–336, 2012.
- [vdBGJ⁺22] Jan van den Brand, Yu Gao, Arun Jambulapati, Yin Tat Lee, Yang P. Liu, Richard Peng, and Aaron Sidford. Faster maxflow via improved dynamic spectral vertex sparsifiers. In Stefano Leonardi and Anupam Gupta, editors, *STOC ’22: 54th Annual ACM SIGACT Symposium on Theory of Computing, Rome, Italy, June 20 - 24, 2022*, pages 543–556. ACM, 2022.
- [vdBLL⁺21] Jan van den Brand, Yin Tat Lee, Yang P. Liu, Thatchaphol Saranurak, Aaron Sidford, Zhao Song, and Di Wang. Minimum cost flows, mdps, and l1-regression in nearly linear time for dense instances. In Samir Khuller and Virginia Vassilevska Williams, editors, *STOC ’21: 53rd Annual ACM SIGACT Symposium on Theory of Computing, Virtual Event, Italy, June 21-25, 2021*, pages 859–869. ACM, 2021.

A SV Approximation Proofs

In this section we provide proofs of equivalences for definitions related to SV-approximation (Appendix A.1), properties of SV-approximation (Appendix A.2), and separations of SV-approximation from other notions of approximation (Appendix A.3).

We recall results useful in the proofs:

Fact A.1 ([CKP⁺17]). Let $\mathbf{A} \in \mathbb{C}^{n \times n}$ be a Hermitian matrix and $I \subset [n]$ be arbitrary. Then \mathbf{A} is PSD if and only if $\mathbf{A}_{I,I}$ is PSD and $\text{SC}_I(\mathbf{A})$ is PSD.

We require a result that matrix approximation is approximately preserved under Schur complements:

Theorem A.2 (Theorem 4.8 [AKM⁺20]). *Given $\mathbf{N}, \tilde{\mathbf{N}} \in \mathbb{C}^{n \times n}$ such that $\|\mathbf{N}\| \leq 1$, suppose $\tilde{\mathbf{N}}$ is an ε -approximation of \mathbf{N} with respect to $\mathbf{S}_{\mathbf{N}}$. Then for $F \subset [n]$ such that $\mathbf{I}_{F,F} - \mathbf{N}_{F,F}$ is invertible, we have that $\mathbf{I}_{F^c} - \mathbf{S}_{\mathbf{C}_F}(\mathbf{I} - \tilde{\mathbf{N}})$ is an $\varepsilon + O(\varepsilon^2)$ approximation of $\mathbf{I}_{F^c} - \mathbf{S}_{\mathbf{C}_F}(\mathbf{I} - \mathbf{N})$ with respect to $\mathbf{S}_{\mathbf{I}_{F^c} - \mathbf{S}_{\mathbf{C}_F}(\mathbf{I} - \mathbf{N})}$.*

We also prove a useful property of matrix approximation:

Lemma A.3 (Manipulating matrix approximation). *Let $\mathbf{A}, \tilde{\mathbf{A}} \in \mathbb{C}^{m \times n}$, let $\mathbf{E} \in \mathbb{C}^{m \times m}, \mathbf{F} \in \mathbb{C}^{n \times n}$ be PSD and Hermitian, and assume that \mathbf{A} is an ε -approximation of $\tilde{\mathbf{A}}$ with respect to error matrices \mathbf{E} and \mathbf{F} , for some $\varepsilon \geq 0$.*

1. For $\mathbf{U} \in \mathbb{C}^{m' \times m}, \mathbf{V} \in \mathbb{C}^{n \times n'}$, $\mathbf{U}\tilde{\mathbf{A}}\mathbf{V}$ is an ε -approximation of $\mathbf{U}\mathbf{A}\mathbf{V}$ with respect to error matrices \mathbf{E}' and \mathbf{F}' , where $\mathbf{E}' = \mathbf{U}\mathbf{E}\mathbf{U}^*$ and $\mathbf{F}' = \mathbf{V}^*\mathbf{F}\mathbf{V}$.
2. For PSD Hermitian $\mathbf{E}' \in \mathbb{C}^{m \times m}$ and $\mathbf{F}' \in \mathbb{C}^{n \times n}$ with $\mathbf{E} \preceq c\mathbf{E}'$ and $\mathbf{F} \preceq c\mathbf{F}'$ for a constant $c \geq 0$, then $\tilde{\mathbf{A}}$ is a $c\varepsilon$ -approximation of \mathbf{A} with respect to error matrices \mathbf{E}' and \mathbf{F}' .

Proof.

1. For arbitrary $x \in \mathbb{C}^m$ and $y \in \mathbb{C}^n$ the definition of ε -approximation implies,

$$\left| x^*(\mathbf{U}\tilde{\mathbf{A}}\mathbf{V} - \mathbf{U}\mathbf{A}\mathbf{V})y \right| = \left| x^*\mathbf{U}(\tilde{\mathbf{A}} - \mathbf{A})\mathbf{V}y \right| \leq \frac{\varepsilon}{2} (x^*\mathbf{U}\mathbf{E}\mathbf{U}^*x + y^*\mathbf{V}^*\mathbf{F}\mathbf{V}y).$$

2. For arbitrary $x \in \mathbb{C}^m, y \in \mathbb{C}^n$,

$$\left| x^*(\tilde{\mathbf{A}} - \mathbf{A})y \right| \leq \frac{\varepsilon}{2} (x^*\mathbf{E}x + y^*\mathbf{F}y) \leq \frac{c\varepsilon}{2} (x^*\mathbf{E}'x + y^*\mathbf{F}'y). \quad \square$$

A.1 Equivalences

Lemma 3.6 (Conditions for SV approximation to be defined). *Let $\mathbf{A}, \tilde{\mathbf{A}} \in \mathbb{C}^{m \times n}$, and let $\mathbf{D}_{\text{in}} \in \mathbb{C}^{m \times m}$ and $\mathbf{D}_{\text{out}} \in \mathbb{C}^{n \times n}$ be PSD matrices such that $\ker(\mathbf{D}_{\text{in}}) \subseteq \text{lker}(\mathbf{A})$ and $\ker(\mathbf{D}_{\text{out}}) \subseteq \text{rker}(\mathbf{A})$. Then the following are equivalent:*

1. $\sigma_{\max}(\mathbf{D}_{\text{in}}^{+1/2}\mathbf{A}\mathbf{D}_{\text{out}}^{+1/2}) \leq 1$.
2. $\mathbf{D}_{\text{in}} - \mathbf{A}\mathbf{D}_{\text{out}}^+\mathbf{A}^*$ is PSD.
3. $\mathbf{D}_{\text{out}} - \mathbf{A}^*\mathbf{D}_{\text{in}}^+\mathbf{A}$ is PSD.
4. For some scalar $z \in \mathbb{C}$ with $|z| = 1$, $\begin{bmatrix} \mathbf{D}_{\text{in}} & z\mathbf{A} \\ z^*\mathbf{A}^* & \mathbf{D}_{\text{out}} \end{bmatrix}$ is PSD.
5. For every scalar $z \in \mathbb{C}$ with $|z| \leq 1$, $\begin{bmatrix} \mathbf{D}_{\text{in}} & z\mathbf{A} \\ z^*\mathbf{A}^* & \mathbf{D}_{\text{out}} \end{bmatrix}$ is PSD.

Suppose further that $\mathbf{D}_{\text{in}} = \mathbf{diag}(d_{\text{in}})$ and $\mathbf{D}_{\text{out}} = \mathbf{diag}(d_{\text{out}})$ for $d_{\text{in}} \in \mathbb{R}_{\geq 0}^m, d_{\text{out}} \in \mathbb{R}_{\geq 0}^n$. Then Condition 1 below implies Condition 2 below, which implies Conditions 1–5 above.

1. \mathbf{A} is nonnegative, $d_{\text{out}} = \mathbf{A}\vec{1}_n, d_{\text{in}} = \vec{1}_m^T\mathbf{A}$.
2. For all $i \in [n]$, $(d_{\text{out}})_i \geq \sum_j |\mathbf{A}_{i,j}|$, and for all $j \in [m]$, $(d_{\text{in}})_j \geq \sum_i |\mathbf{A}_{i,j}|$.

Proof. We first show Conditions 1 – 5 are equivalent.

1 → 3 Condition 1 implies for all y , $y^* \mathbf{D}_{\text{out}}^{+/2} \mathbf{A}^* \mathbf{D}_{\text{in}}^+ \mathbf{A} \mathbf{D}_{\text{out}}^{+/2} y \leq y^* y$ and letting $y \leftarrow \mathbf{D}_{\text{out}}^{1/2} y$ and using that $\ker(\mathbf{D}_{\text{out}}) \subseteq \ker(\mathbf{A})$ implies Condition 3.

3 → 1 Condition 3 implies for all y we have $y^* \mathbf{A}^* \mathbf{D}_{\text{in}}^+ \mathbf{A} y \leq y^* \mathbf{D}_{\text{out}} y$ and taking $y \leftarrow \mathbf{D}_{\text{out}}^{+/2} y$ we have $y^* \mathbf{D}_{\text{out}}^{+/2} \mathbf{A}^* \mathbf{D}_{\text{in}}^+ \mathbf{A} \mathbf{D}_{\text{out}}^{+/2} y \leq y^* \mathbf{D}_{\text{out}} y \leq y^* y$ which implies Condition 1.

3 → 5 Letting z with $|z| \leq 1$ be arbitrary, we have that by Condition 3 $\mathbf{D}_{\text{in}} - z^* \mathbf{A}^* \mathbf{D}_{\text{out}}^+ z \mathbf{A} = \mathbf{D}_{\text{in}} - \mathbf{A}^* \mathbf{D}_{\text{out}}^+ \mathbf{A} \succeq \mathbf{0}$, and since $\mathbf{D}_{\text{in}} \succeq \mathbf{0}$ we apply Fact A.1 and conclude Condition 5. An analogous argument shows 2 → 5. The fact that Fact A.1 is an equivalence then implies 5 → 2 and 5 → 3.

5 → 4 Immediate.

4 → 2 By Fact A.1 and the fact that $z^* z = 1$ we have $\mathbf{D}_{\text{in}} - z^* \mathbf{A}^* \mathbf{D}_{\text{out}}^+ z \mathbf{A} = \mathbf{D}_{\text{in}} - \mathbf{A}^* \mathbf{D}_{\text{out}}^+ \mathbf{A} \succeq \mathbf{0}$.

We next show the latter set of conditions.

1 → 2 For all $i \in [n]$ we have $[d_{\text{out}}]_i = \sum_j \mathbf{A}_{i,j} = \sum_j |\mathbf{A}_{i,j}|$.

2 → 5 We use that a Hermitian matrix that is diagonally dominant is positive semidefinite. We have $(\mathbf{D}_{\text{in}})_{i,i} \geq \sum_j |z \mathbf{A}_{i,j}|$ and $(\mathbf{D}_{\text{out}})_{i,i} \geq \sum_j |z^* \mathbf{A}_{j,i}^*|$ so $\begin{bmatrix} \mathbf{D}_{\text{in}} & \mathbf{A} \\ \mathbf{A}^* & \mathbf{D}_{\text{out}} \end{bmatrix}$ is diagonally dominant and thus PSD. \square

Lemma 3.7 (Equivalent formulations of SV approximation). *Let $\mathbf{A}, \tilde{\mathbf{A}} \in \mathbb{C}^{m \times n}$ and let $\mathbf{D}_{\text{in}} \in \mathbb{C}^{m \times m}$ and $\mathbf{D}_{\text{out}} \in \mathbb{C}^{n \times n}$ be PSD matrices. Then the following are equivalent*

1. $\tilde{\mathbf{A}} \stackrel{\mathbf{D}_{\text{in}}, \mathbf{D}_{\text{out}}}{\approx_{\varepsilon}} \mathbf{A}$.

2. $\text{sift}(\tilde{\mathbf{A}}) \stackrel{\mathbf{D}}{\approx_{\varepsilon}} \text{sift}(\tilde{\mathbf{B}})$, where

$$\text{sift}(\tilde{\mathbf{A}}) = \begin{bmatrix} \mathbf{0}^{m \times m} & \tilde{\mathbf{A}} \\ \tilde{\mathbf{A}}^* & \mathbf{0}^{n \times n} \end{bmatrix}, \text{sift}(\mathbf{A}) = \begin{bmatrix} \mathbf{0}^{m \times m} & \mathbf{A} \\ \mathbf{A}^* & \mathbf{0}^{n \times n} \end{bmatrix}, \mathbf{D} = \begin{bmatrix} \mathbf{D}_{\text{in}} & \mathbf{0}^{m \times n} \\ \mathbf{0}^{n \times m} & \mathbf{D}_{\text{out}} \end{bmatrix}.$$

3. For some scalar $z \in \mathbb{C}$ with $|z| = 1$, $\tilde{\mathbf{C}}$ is $\varepsilon/2$ -approximation of \mathbf{C} with respect to error matrix \mathbf{E} , where

$$\mathbf{C} = \begin{bmatrix} \mathbf{0}^{m \times m} & z \mathbf{A} \\ \mathbf{0}^{n \times m} & \mathbf{0}^{n \times n} \end{bmatrix}, \tilde{\mathbf{C}} = \begin{bmatrix} \mathbf{0}^{m \times m} & z \tilde{\mathbf{A}} \\ \mathbf{0}^{n \times m} & \mathbf{0}^{n \times n} \end{bmatrix}, \mathbf{E} = \begin{bmatrix} \mathbf{D}_{\text{in}} & z \mathbf{A} \\ z^* \mathbf{A}^* & \mathbf{D}_{\text{out}} \end{bmatrix}$$

4. Item 3 holds for every $z \in \mathbb{C}$ such that $|z| \leq 1$.

Proof. We prove 1 \iff 2 and 1 → 4 → 3 → 1.

1 \iff 2 Suppose Item 1 holds. Then for arbitrary test vectors $x = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$, $y = \begin{bmatrix} y_1 \\ y_2 \end{bmatrix}$ where $x_1, y_1 \in \mathbb{C}^m$ and $x_2, y_2 \in \mathbb{C}^n$ we have

$$\begin{aligned} \left| x^* (\mathbf{B} - \tilde{\mathbf{B}}) y \right| &\leq \left| x_1^* (\mathbf{A} - \tilde{\mathbf{A}}) y_2 \right| + \left| x_2^* (\mathbf{A}^* - \tilde{\mathbf{A}}^*) y_1 \right| \\ &\leq \frac{\varepsilon}{4} (x_1^* \mathbf{D}_{\text{in}} x_1 + y_2^* \mathbf{D}_{\text{out}} y_2 + x_1^* \mathbf{A} \mathbf{D}_{\text{out}}^+ \mathbf{A}^* x_1 + y_2^* \mathbf{A}^* \mathbf{D}_{\text{in}}^+ \mathbf{A} y_2) \\ &\quad + \frac{\varepsilon}{4} (x_2^* \mathbf{D}_{\text{out}} x_2 + y_1^* \mathbf{D}_{\text{in}} y_1 + x_2^* \mathbf{A}^* \mathbf{D}_{\text{in}}^+ \mathbf{A} x_2 + y_1^* \mathbf{A} \mathbf{D}_{\text{out}}^+ \mathbf{A}^* y_1) \\ &= \frac{\varepsilon}{4} (x^* \mathbf{D} x + y^* \mathbf{D} y + x^* \mathbf{B} \mathbf{D}^{-1} \mathbf{B}^* x + y^* \mathbf{B}^* \mathbf{D}^{-1} \mathbf{B} y). \end{aligned}$$

Furthermore, $\ker(\mathbf{D}) \subseteq \text{lker}(\mathbf{B}), \ker(\mathbf{D}) \subseteq \text{rker}(\mathbf{B})$ and $\mathbf{D} - \mathbf{B}\mathbf{D}^+\mathbf{B} \succeq \mathbf{0}$ by Item 5 of Lemma 3.6. In the other direction, we obtain $2 \rightarrow 1$ by considering the set of test vectors $\begin{bmatrix} x_1 \\ 0^n \end{bmatrix}$ and $\begin{bmatrix} 0^m \\ y_2 \end{bmatrix}$.

1 \rightarrow 4: Fix arbitrary z with $|z| \leq 1$. We have that $\mathbf{E} \succeq \mathbf{0}$ by Item 5 of Lemma 3.6. Then for arbitrary test vectors $x = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}, y = \begin{bmatrix} y_1 \\ y_2 \end{bmatrix}$, let $x' = \begin{bmatrix} x_1 \\ x'_2 \end{bmatrix}, y' = \begin{bmatrix} y_1 \\ y'_2 \end{bmatrix}$ where

$$x'_2 = \operatorname{argmin}_{v \in \mathbb{C}^n} \begin{bmatrix} x_1^* & v^* \end{bmatrix} \begin{bmatrix} \mathbf{D}_{\text{in}} & z\mathbf{A} \\ z^*\mathbf{A}^* & \mathbf{D}_{\text{out}} \end{bmatrix} \begin{bmatrix} x_1 \\ v \end{bmatrix}, \quad y'_2 = \operatorname{argmin}_{v \in \mathbb{C}^m} \begin{bmatrix} y_2^* & v^* \end{bmatrix} \begin{bmatrix} \mathbf{D}_{\text{out}} & z^*\mathbf{A}^* \\ z\mathbf{A} & \mathbf{D}_{\text{in}} \end{bmatrix} \begin{bmatrix} y_2 \\ v \end{bmatrix}.$$

Thus

$$x'^* \begin{bmatrix} \mathbf{D}_{\text{in}} & z\mathbf{A} \\ z^*\mathbf{A}^* & \mathbf{D}_{\text{out}} \end{bmatrix} x' = x_1^* \text{SC} \left(\begin{bmatrix} \mathbf{D}_{\text{in}} & z\mathbf{A} \\ z^*\mathbf{A}^* & \mathbf{D}_{\text{out}} \end{bmatrix} \right) x_1, \quad y'^* \begin{bmatrix} \mathbf{D}_{\text{in}} & z\mathbf{A} \\ z^*\mathbf{A}^* & \mathbf{D}_{\text{out}} \end{bmatrix} y' = y_2^* \text{SC} \left(\begin{bmatrix} \mathbf{D}_{\text{out}} & z^*\mathbf{A}^* \\ z\mathbf{A} & \mathbf{D}_{\text{in}} \end{bmatrix} \right) y_2.$$

Then we have

$$\begin{aligned} \left| x^*(\tilde{\mathbf{C}} - \mathbf{C})y \right| &= \left| (x_1 z^*)^*(\tilde{\mathbf{A}} - \mathbf{A})(y_2 z) \right| \\ &\leq \frac{\varepsilon}{4} (x_1^*(\mathbf{D}_{\text{in}} - z\mathbf{A}\mathbf{D}_{\text{out}}^+\mathbf{A}^*z^*)x_1 + y_2^*(\mathbf{D}_{\text{out}} - z^*\mathbf{A}^*\mathbf{D}_{\text{in}}^+\mathbf{A}z)y_2) \\ &= \frac{\varepsilon}{4} \left(x_1^* \text{SC} \left(\begin{bmatrix} \mathbf{D}_{\text{in}} & z\mathbf{A} \\ z^*\mathbf{A}^* & \mathbf{D}_{\text{out}} \end{bmatrix} \right) x_1 + y_2^* \text{SC} \left(\begin{bmatrix} \mathbf{D}_{\text{out}} & z\mathbf{A} \\ z^*\mathbf{A}^* & \mathbf{D}_{\text{in}} \end{bmatrix} \right) y_2 \right) \\ &= \frac{\varepsilon}{4} \left(x'^* \begin{bmatrix} \mathbf{D}_{\text{in}} & z\mathbf{A} \\ z^*\mathbf{A}^* & \mathbf{D}_{\text{out}} \end{bmatrix} x' + y'^* \begin{bmatrix} \mathbf{D}_{\text{in}} & z\mathbf{A} \\ z^*\mathbf{A}^* & \mathbf{D}_{\text{out}} \end{bmatrix} y' \right) \\ &\leq \frac{\varepsilon}{4} (x^*\mathbf{E}x + y^*\mathbf{E}y). \end{aligned}$$

4 \rightarrow 3 Immediate.

3 \rightarrow 1 For arbitrary test vectors $x_1 \in \mathbb{C}^m, y_2 \in \mathbb{C}^n$, let $x = \begin{bmatrix} zx_1 \\ x_2 \end{bmatrix}, y = \begin{bmatrix} y_1 \\ z^*y_2 \end{bmatrix}$ where

$$x_2 = \operatorname{argmin}_{v \in \mathbb{C}^n} \begin{bmatrix} z^*x_1^* & v^* \end{bmatrix} \begin{bmatrix} \mathbf{D}_{\text{in}} & z\mathbf{A} \\ z^*\mathbf{A}^* & \mathbf{D}_{\text{out}} \end{bmatrix} \begin{bmatrix} zx_1 \\ v \end{bmatrix}, \quad y_1 = \operatorname{argmin}_{v \in \mathbb{C}^m} \begin{bmatrix} zy_2^* & v^* \end{bmatrix} \begin{bmatrix} \mathbf{D}_{\text{out}} & z^*\mathbf{A}^* \\ z\mathbf{A} & \mathbf{D}_{\text{in}} \end{bmatrix} \begin{bmatrix} z^*y_2 \\ v \end{bmatrix}$$

Then

$$\begin{aligned} \left| x_1^*(\tilde{\mathbf{A}} - \mathbf{A})y_2 \right| &= \left| x^*(\tilde{\mathbf{C}} - \mathbf{C})x \right| \\ &\leq \frac{\varepsilon}{4} (x^*\mathbf{E}x + y^*\mathbf{E}y) \\ &= \frac{\varepsilon}{4} \left(z^*x_1^* \text{SC} \left(\begin{bmatrix} \mathbf{D}_{\text{in}} & z\mathbf{A} \\ z^*\mathbf{A}^* & \mathbf{D}_{\text{out}} \end{bmatrix} \right) x_1 z + zy_2^* \text{SC} \left(\begin{bmatrix} \mathbf{D}_{\text{out}} & z\mathbf{A} \\ z^*\mathbf{A}^* & \mathbf{D}_{\text{in}} \end{bmatrix} \right) z^*y_2 \right) \\ &= \frac{\varepsilon}{4} (z^*x_1^*(\mathbf{D}_{\text{in}} - z\mathbf{A}\mathbf{D}_{\text{out}}^+\mathbf{A}^*z^*)zx_1 + zy_2^*(\mathbf{D}_{\text{out}} - z^*\mathbf{A}^*\mathbf{D}_{\text{in}}^+\mathbf{A}z)z^*y_2) \\ &= \frac{\varepsilon}{4} (x_1^*(\mathbf{D}_{\text{in}} - \mathbf{A}\mathbf{D}_{\text{out}}^+\mathbf{A}^*)x_1 + y_2^*(\mathbf{D}_{\text{out}} - \mathbf{A}^*\mathbf{D}_{\text{in}}^+\mathbf{A})y_2). \quad \square \end{aligned}$$

Lemma 3.8. *Let $\mathbf{A}, \tilde{\mathbf{A}} \in \mathbb{C}^{m \times n}$, and let $\mathbf{D}_{\text{in}} \in \mathbb{C}^{m \times m}$ and $\mathbf{D}_{\text{out}} \in \mathbb{C}^{n \times n}$ be PSD matrices such that $\ker(\mathbf{D}_{\text{in}}) \subseteq \text{lker}(\mathbf{A})$, and $\ker(\mathbf{D}_{\text{out}}) \subseteq \text{rker}(\mathbf{A})$. Let $\mathbf{N} = \mathbf{D}_{\text{in}}^{+1/2} \mathbf{A} \mathbf{D}_{\text{out}}^{+1/2}$ and $\tilde{\mathbf{N}} = \mathbf{D}_{\text{in}}^{+1/2} \tilde{\mathbf{A}} \mathbf{D}_{\text{out}}^{+1/2}$. Then $\tilde{\mathbf{A}} \stackrel{\text{D}_{\text{in}}, \text{D}_{\text{out}}}{\approx_{\varepsilon}} \mathbf{A}$ if and only if $\tilde{\mathbf{N}} \stackrel{\text{svn}}{\approx_{\varepsilon}} \mathbf{N}$.*

Proof. Suppose $\tilde{\mathbf{A}} \stackrel{\text{D}_{\text{in}}, \text{D}_{\text{out}}}{\approx_{\varepsilon}} \mathbf{A}$, i.e. $\tilde{\mathbf{A}}$ is an $\varepsilon/2$ -approximation of \mathbf{A} with respect to

$$\mathbf{E} = \mathbf{D}_{\text{in}} - \mathbf{A} \mathbf{D}_{\text{out}}^+ \mathbf{A}^*, \quad \mathbf{F} = \mathbf{D}_{\text{out}} - \mathbf{A} \mathbf{D}_{\text{in}}^+ \mathbf{A}^*.$$

Applying Lemma A.3 with $\mathbf{U} = \mathbf{D}_{\text{in}}^{+1/2}$, $\mathbf{V} = \mathbf{D}_{\text{out}}^{+1/2}$ we obtain that $\tilde{\mathbf{N}}$ is an $\varepsilon/2$ -approximation of \mathbf{N} with respect to error matrices

$$\mathbf{D}_{\text{in}}^{+1/2} (\mathbf{D}_{\text{in}} - \mathbf{A} \mathbf{D}_{\text{out}}^+ \mathbf{A}^*) \mathbf{D}_{\text{in}}^{+1/2} \preceq \mathbf{I} - (\mathbf{D}_{\text{in}}^{+1/2} \mathbf{A} \mathbf{D}_{\text{out}}^{+1/2}) (\mathbf{D}_{\text{out}}^{+1/2} \mathbf{A}^* \mathbf{D}_{\text{in}}^{+1/2}) = \mathbf{I} - \mathbf{N} \mathbf{N}^*$$

and

$$\mathbf{D}_{\text{out}}^{+1/2} (\mathbf{D}_{\text{out}} - \mathbf{A}^* \mathbf{D}_{\text{in}}^+ \mathbf{A}) \mathbf{D}_{\text{out}}^{+1/2} \preceq \mathbf{I} - (\mathbf{D}_{\text{out}}^{+1/2} \mathbf{A}^* \mathbf{D}_{\text{in}}^{+1/2}) (\mathbf{D}_{\text{in}}^{+1/2} \mathbf{A} \mathbf{D}_{\text{out}}^{+1/2}) = \mathbf{I} - \mathbf{N}^* \mathbf{N}$$

and hence we obtain that $\tilde{\mathbf{N}} \stackrel{\text{svn}}{\approx_{\varepsilon}} \mathbf{N}$. The other direction is analogous, and the kernel properties follow as $\ker(\mathbf{D}_{\text{in}}) \subseteq \text{lker}(\mathbf{D}_{\text{in}}^{1/2} \mathbf{D}_{\text{in}}^{+1/2} \mathbf{A} \mathbf{D}_{\text{out}}^{+1/2} \mathbf{D}_{\text{out}}^{1/2})$ and $\ker(\mathbf{D}_{\text{out}}) \subseteq \text{rker}(\mathbf{D}_{\text{in}}^{1/2} \mathbf{D}_{\text{in}}^{+1/2} \mathbf{A} \mathbf{D}_{\text{out}}^{+1/2} \mathbf{D}_{\text{out}}^{1/2})$. \square

Theorem 3.9 (Unitary transformation characterization of SV-approximation). *For $\tilde{\mathbf{N}}, \mathbf{N} \in \mathbb{C}^{n \times n}$, we have that $\tilde{\mathbf{N}} \stackrel{\text{svn}}{\approx_{\varepsilon}} \mathbf{N}$ if for every pair of unitary matrices \mathbf{U}, \mathbf{V} , $\mathbf{U} \tilde{\mathbf{N}} \mathbf{V}$ is a standard $\varepsilon/2$ -approximation of $\mathbf{U} \mathbf{N} \mathbf{V}$ with respect to degree matrix \mathbf{I} . Moreover, if $\tilde{\mathbf{N}} \stackrel{\text{svn}}{\approx_{\varepsilon}} \mathbf{N}$ then for every pair of matrices \mathbf{U}, \mathbf{V} satisfying $\|\mathbf{U}\| \leq 1, \|\mathbf{V}\| \leq 1$, we have that $\mathbf{U} \tilde{\mathbf{N}} \mathbf{V} \stackrel{\text{svn}}{\approx_{\varepsilon}} \mathbf{U} \mathbf{N} \mathbf{V}$, and hence $\mathbf{U} \tilde{\mathbf{N}} \mathbf{V}$ is a standard ε -approximation of $\mathbf{U} \mathbf{N} \mathbf{V}$ with respect to degree matrix \mathbf{I} .*

Proof.

- Suppose for every pair of unitary matrices \mathbf{U}, \mathbf{V} , we have for all $x, y \in \mathbb{C}^n$ that

$$|x^* (\mathbf{U} \tilde{\mathbf{A}} \mathbf{V} - \mathbf{U} \mathbf{A} \mathbf{V}) y| \leq \frac{\varepsilon}{2} \cdot \sqrt{x^* (\mathbf{I} - \mathbf{S}_{\mathbf{U} \mathbf{A} \mathbf{V}}) x} \cdot \sqrt{y^* (\mathbf{I} - \mathbf{S}_{\mathbf{U} \mathbf{A} \mathbf{V}}) y}.$$

We will pick $u, v, \mathbf{U}, \mathbf{V}$ to depend on x, y and the result will follow immediately from invoking the above equation with $x, y \leftarrow u, v$.

Set $\mathbf{U} = \mathbf{I}$, $u = x/\|x\|$, and $v = \mathbf{A}y/\|\mathbf{A}y\|$. Note that $u^*v = (x^* \mathbf{A}y)/(\|x\| \cdot \|y\|)$, so u and v have the same angle between them as \mathbf{A}^*x and y . Hence, there exists a unitary matrix \mathbf{V} such that $\mathbf{V}u = \mathbf{A}^*x/\|\mathbf{A}^*x\|$ and $\mathbf{V}v = y/\|y\|$. Substituting gives

$$\frac{|x^* (\tilde{\mathbf{A}} - \mathbf{A}) y|}{\|x\| \|y\|} \leq \frac{\varepsilon}{2} \cdot \sqrt{1 - \frac{x^* (\mathbf{A} \mathbf{A}^*) x}{\|x\| \|\mathbf{A}^* x\|}} \cdot \sqrt{1 - \frac{y^* (\mathbf{A}^* \mathbf{A}) y}{\|y\| \|\mathbf{A} y\|}}.$$

Applying $\|\mathbf{A}^*\| = \|\mathbf{A}\| \leq 1$ gives the desired result.

- In the other direction, we start with Item 3 of Lemma 3.7 with $z = 1$ applied to the approximation statement $\tilde{\mathbf{N}} \stackrel{\text{svn}}{\approx_{\varepsilon}} \mathbf{N}$ and apply Lemma A.3 with left and right hand side matrices

$$\mathbf{U}' \leftarrow \begin{bmatrix} \mathbf{U} & \mathbf{0} \\ \mathbf{0} & \mathbf{V}^* \end{bmatrix} \quad \text{and} \quad \mathbf{V}' \leftarrow \begin{bmatrix} \mathbf{U}^* & \mathbf{0} \\ \mathbf{0} & \mathbf{V} \end{bmatrix}$$

This results in the approximation statement that for the matrices given below, $\mathbf{R}, \tilde{\mathbf{R}} \epsilon/2$ -approximate each other with respect to \mathbf{E} where

$$\mathbf{R} = \begin{bmatrix} \mathbf{0} & \mathbf{UNV} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \quad \tilde{\mathbf{R}} = \begin{bmatrix} \mathbf{0} & \mathbf{U\tilde{N}V} \\ \mathbf{0} & \mathbf{0} \end{bmatrix}, \text{ and} \quad \mathbf{E} = \begin{bmatrix} \mathbf{UIU}^* & \mathbf{UNV} \\ (\mathbf{UNV})^* & \mathbf{V}^*\mathbf{IV} \end{bmatrix} \preceq \begin{bmatrix} \mathbf{I} & \mathbf{UNV} \\ (\mathbf{UNV})^* & \mathbf{I} \end{bmatrix}.$$

Hence, by Item 3 of Lemma 3.7 we conclude $\mathbf{UNV} \overset{svn}{\approx}_{\epsilon} \mathbf{U\tilde{N}V}$. \square

Corollary 3.10 (SV preservation under multiplication by permutation matrices). *Let $\mathbf{A}, \tilde{\mathbf{A}} \in \mathbb{C}_{\geq 0}^{m \times n}$ and suppose $\tilde{\mathbf{A}} \overset{\mathbf{D}_{\text{in}}, \mathbf{D}_{\text{out}}}{\approx}_{\epsilon} \mathbf{A}$. Let \mathbf{U}, \mathbf{V} be arbitrary permutation matrices. Then $\mathbf{U}\tilde{\mathbf{A}}\mathbf{V} \overset{\mathbf{D}'_{\text{in}}, \mathbf{D}'_{\text{out}}}{\approx}_{\epsilon} \mathbf{UAV}$ where $\mathbf{D}'_{\text{in}} = \mathbf{UD}_{\text{in}}\mathbf{U}^*$ and $\mathbf{D}'_{\text{out}} = \mathbf{V}^*\mathbf{D}_{\text{out}}\mathbf{V}$. Consequently, if $\tilde{\mathbf{A}} \overset{sv}{\approx}_{\epsilon} \mathbf{A}$, then $\mathbf{U}\tilde{\mathbf{A}}\mathbf{V} \overset{sv}{\approx}_{\epsilon} \mathbf{UAV}$ and if $\tilde{\mathbf{A}} \overset{svn}{\approx}_{\epsilon} \mathbf{A}$ then $\mathbf{U}\tilde{\mathbf{A}}\mathbf{V} \overset{svn}{\approx}_{\epsilon} \mathbf{UAV}$.*

We start with Item 3 of Lemma 3.7 with $z = 1$ applied to the approximation statement we are given in the lemma statement and apply Lemma A.3 with left and right hand side matrices

$$\mathbf{U}' \leftarrow \begin{bmatrix} \mathbf{U} & \mathbf{0} \\ \mathbf{0} & \mathbf{V}^* \end{bmatrix} \quad \text{and} \quad \mathbf{V}' \leftarrow \begin{bmatrix} \mathbf{U}^* & \mathbf{0} \\ \mathbf{0} & \mathbf{V} \end{bmatrix}$$

This results in the approximation statement that for the matrices given below, $\mathbf{R}, \tilde{\mathbf{R}} \epsilon/2$ -approximate each other with respect to \mathbf{E} where

$$\mathbf{R} = \begin{bmatrix} \mathbf{0} & \mathbf{UAV} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \quad \tilde{\mathbf{R}} = \begin{bmatrix} \mathbf{0} & \mathbf{U\tilde{A}V} \\ \mathbf{0} & \mathbf{0} \end{bmatrix}, \text{ and} \quad \mathbf{E} = \begin{bmatrix} \mathbf{UD}_{\text{in}}\mathbf{U}^* & \mathbf{UAV} \\ (\mathbf{UAV})^* & \mathbf{V}^*\mathbf{D}_{\text{out}}\mathbf{V} \end{bmatrix}.$$

Hence, Item 3 of Lemma 3.7 is satisfied for SV approximation of \mathbf{UAV} and $\mathbf{U\tilde{A}V}$ with respect to $\mathbf{UD}_{\text{in}}\mathbf{U}^*$ and $\mathbf{V}^*\mathbf{D}_{\text{out}}\mathbf{V}$.

For the first ‘‘consequently’’ claim, observe that $\mathbf{UAV}\vec{\mathbf{1}} = \mathbf{U}\vec{\mathbf{1}}$ since \mathbf{V} is a permutation. Hence,

$$\text{diag}(\mathbf{UAV}\vec{\mathbf{1}}) = \mathbf{UD}_{\text{in}}\mathbf{U}^*.$$

A similar analysis shows

$$\text{diag}(\vec{\mathbf{1}}^{\top} \mathbf{UAV}) = \mathbf{V}^*\mathbf{D}_{\text{in}}\mathbf{V}.$$

the second statement follows from $\mathbf{UU}^* = \mathbf{V}^*\mathbf{V} = \mathbf{I}$.

A.2 Properties

We prove that SV approximation implies standard approximation:

Lemma A.4. *Let $\mathbf{A}, \tilde{\mathbf{A}} \in \mathbb{C}^{n \times n}$ and suppose $\tilde{\mathbf{A}} \overset{\mathbf{D}_{\text{in}}, \mathbf{D}_{\text{out}}}{\approx}_{\epsilon} \mathbf{A}$. Then, $\tilde{\mathbf{A}}$ is an ϵ -approximation of \mathbf{A} with respect to $\mathbf{E} = \mathbf{D}_{\text{in}} - \mathbf{A}$, $\mathbf{F} = \mathbf{D}_{\text{out}} - \mathbf{A}^*$.*

Proof. This follows immediately from specializing Item 3 of Lemma 3.7 with $z = -1$ to test vectors of the form $\begin{bmatrix} x \\ x \end{bmatrix}, \begin{bmatrix} y \\ y \end{bmatrix}$. \square

It likewise implies UC approximation:

Lemma 3.11. *If $\mathbf{A}, \tilde{\mathbf{A}} \in \mathbb{C}^{n \times n}$ with $\tilde{\mathbf{A}} \overset{\mathbf{D}}{\approx}_{\epsilon} \mathbf{A}$ then $\tilde{\mathbf{A}} \overset{\circ}{\approx}_{\epsilon} \mathbf{A}$ with respect to degree matrix \mathbf{D} .*

Proof. By Item 4 of Lemma 3.7 we have that for every pair of test vectors $\begin{bmatrix} x \\ x \end{bmatrix}, \begin{bmatrix} y \\ y \end{bmatrix}$ and every unit magnitude z , we have

$$\left| x^*(z\mathbf{A} - z\tilde{\mathbf{A}})y \right| \leq \frac{\varepsilon}{4} \cdot 2(x^*\mathbf{D}x + y^*\mathbf{D}y + \Re(zx^*\mathbf{A}x + zy\mathbf{A}y^*))$$

Then choosing z to minimize $\Re(zx^*\mathbf{A}x + zy\mathbf{A}y^*)$, we obtain

$$\left| x^*(\mathbf{A} - \tilde{\mathbf{A}})y \right| \leq \frac{2\varepsilon}{4} (x^*\mathbf{D}x + y^*\mathbf{D}y - |x^*\mathbf{A}x + y\mathbf{A}y^*|)$$

Which is precisely the condition for $\tilde{\mathbf{A}} \overset{\circ}{\approx}_{\varepsilon} \mathbf{A}$ with respect to \mathbf{D} . \square

Lemma 3.14 (SV preservation under arbitrary lifting). *Let $\mathbf{A}, \tilde{\mathbf{A}} \in \mathbb{C}^{m \times n}$ be matrices such that $\tilde{\mathbf{A}} \overset{\mathbf{D}_{\text{in}}, \mathbf{D}_{\text{out}}}{\approx}_{\varepsilon} \mathbf{A}$. Then for all integers $i, j, k, \ell \geq 0$*

$$\begin{bmatrix} \mathbf{0}^{i \times j} & \mathbf{0}^{i \times n} & \mathbf{0}^{i \times k} \\ \mathbf{0}^{m \times j} & \tilde{\mathbf{A}} & \mathbf{0}^{m \times k} \\ \mathbf{0}^{\ell \times j} & \mathbf{0}^{\ell \times n} & \mathbf{0}^{\ell \times k} \end{bmatrix} \overset{\mathbf{D}'_{\text{in}}, \mathbf{D}'_{\text{out}}}{\approx}_{\varepsilon} \begin{bmatrix} \mathbf{0}^{i \times j} & \mathbf{0}^{i \times n} & \mathbf{0}^{i \times k} \\ \mathbf{0}^{m \times j} & \mathbf{A} & \mathbf{0}^{m \times k} \\ \mathbf{0}^{\ell \times j} & \mathbf{0}^{\ell \times n} & \mathbf{0}^{\ell \times k} \end{bmatrix}.$$

Where

$$\mathbf{D}'_{\text{in}} = \begin{bmatrix} \mathbf{0}^{i \times i} & \mathbf{0}^{i \times n} & \mathbf{0}^{i \times \ell} \\ \mathbf{0}^{n \times i} & \mathbf{D}_{\text{in}} & \mathbf{0}^{n \times \ell} \\ \mathbf{0}^{\ell \times i} & \mathbf{0}^{\ell \times n} & \mathbf{0}^{\ell \times \ell} \end{bmatrix}, \quad \mathbf{D}'_{\text{out}} = \begin{bmatrix} \mathbf{0}^{j \times j} & \mathbf{0}^{j \times m} & \mathbf{0}^{j \times k} \\ \mathbf{0}^{m \times j} & \mathbf{D}_{\text{out}} & \mathbf{0}^{m \times k} \\ \mathbf{0}^{k \times j} & \mathbf{0}^{k \times m} & \mathbf{0}^{k \times k} \end{bmatrix}.$$

Consequently, if $\tilde{\mathbf{A}} \overset{sv}{\approx}_{\varepsilon} \mathbf{A}$ then

$$\begin{bmatrix} \mathbf{0}^{i \times j} & \mathbf{0}^{i \times n} & \mathbf{0}^{i \times k} \\ \mathbf{0}^{m \times j} & \tilde{\mathbf{A}} & \mathbf{0}^{m \times k} \\ \mathbf{0}^{\ell \times j} & \mathbf{0}^{\ell \times n} & \mathbf{0}^{\ell \times k} \end{bmatrix} \overset{sv}{\approx}_{\varepsilon} \begin{bmatrix} \mathbf{0}^{i \times j} & \mathbf{0}^{i \times n} & \mathbf{0}^{i \times k} \\ \mathbf{0}^{m \times j} & \mathbf{A} & \mathbf{0}^{m \times k} \\ \mathbf{0}^{\ell \times j} & \mathbf{0}^{\ell \times n} & \mathbf{0}^{\ell \times k} \end{bmatrix}.$$

Proof. Observe by expanding the definition of SV approximation that it is preserved under embedding into the first principal diagonal block of any larger matrix that is zeros elsewhere. That is,

$$\begin{bmatrix} \tilde{\mathbf{A}} & \mathbf{0}^{m \times (j+k)} \\ \mathbf{0}^{(i+\ell) \times n} & \mathbf{0}^{(i+\ell) \times (j+k)} \end{bmatrix} \overset{\mathbf{D}'_{\text{in}}, \mathbf{D}'_{\text{out}}}{\approx}_{\varepsilon} \begin{bmatrix} \mathbf{A} & \mathbf{0}^{m \times (j+k)} \\ \mathbf{0}^{(i+\ell) \times n} & \mathbf{0}^{(i+\ell) \times (j+k)} \end{bmatrix} \quad (29)$$

where $\mathbf{D}'_{\text{in}}, \mathbf{D}'_{\text{out}}$ are the respective principle embeddings of the degree matrices.

Let \mathbf{U}, \mathbf{V} be permutation matrices such that

$$\mathbf{U} \begin{bmatrix} \mathbf{A} & \mathbf{0}^{m \times (j+k)} \\ \mathbf{0}^{(i+\ell) \times n} & \mathbf{0}^{(i+\ell) \times (j+k)} \end{bmatrix} \mathbf{V} = \begin{bmatrix} \mathbf{0}^{i \times j} & \mathbf{0}^{i \times n} & \mathbf{0}^{i \times k} \\ \mathbf{0}^{m \times j} & \mathbf{A} & \mathbf{0}^{m \times k} \\ \mathbf{0}^{\ell \times j} & \mathbf{0}^{\ell \times n} & \mathbf{0}^{\ell \times k} \end{bmatrix}.$$

It suffices to argue that we can apply this transformation to both sides of Equation (29) and have the approximation still hold. This is implied by Corollary 3.10 with $\mathbf{D}'_{\text{in}} \leftarrow \mathbf{U}\mathbf{D}'_{\text{in}}\mathbf{U}^*$ and $\mathbf{D}'_{\text{out}} \leftarrow \mathbf{V}^*\mathbf{D}'_{\text{out}}\mathbf{V}$. The ‘‘consequently’’ claim holds from inspecting the resulting block structure. \square

Lemma 3.16 (SV preservation under products). *Let $(\mathbf{N}_i)_{i \in [\ell]}, (\tilde{\mathbf{N}}_i)_{i \in [\ell]} \in \mathbb{C}^{n \times n}$ be such that for every i , $\tilde{\mathbf{N}}_i \overset{svn}{\approx}_{\varepsilon} \mathbf{N}_i$. Then $\tilde{\mathbf{N}}_{\ell} \cdots \tilde{\mathbf{N}}_2 \tilde{\mathbf{N}}_1 \overset{svn}{\approx}_{\varepsilon + O(\varepsilon^2)} \mathbf{N}_{\ell} \cdots \mathbf{N}_2 \mathbf{N}_1$.*

Proof. Using preservation under lifting and sums (Lemmas 3.14 and 3.18) and that SV approximation implies standard approximation for square matrices (Lemma A.4), we can obtain the approximation statement that the matrices $\mathbf{R}, \tilde{\mathbf{R}}$ given below $\epsilon/2$ -standard-approximate each other with respect to the symmetrization $\mathbf{S}_{\mathbf{R}}$ of the first one:

$$\mathbf{R} = \begin{bmatrix} \mathbf{I}/2 & \mathbf{N}_\ell & & \mathbf{0} \\ \mathbf{0} & \mathbf{I} & \mathbf{N}_{\ell-1} & \vdots \\ \vdots & & \ddots & \ddots \\ \vdots & & & \mathbf{I} & \mathbf{N}_1 \\ \mathbf{0}^{n \times n} & & & \mathbf{0} & \mathbf{I}/2 \end{bmatrix} \quad \text{and} \quad \tilde{\mathbf{R}} = \begin{bmatrix} \mathbf{I}/2 & \tilde{\mathbf{N}}_\ell & & \mathbf{0} \\ \mathbf{0} & \mathbf{I} & \tilde{\mathbf{N}}_{\ell-1} & \vdots \\ \vdots & & \ddots & \ddots \\ \vdots & & & \mathbf{I} & \tilde{\mathbf{N}}_1 \\ \mathbf{0}^{n \times n} & & & \mathbf{0} & \mathbf{I}/2 \end{bmatrix}.$$

Taking the Schur complement by eliminating the center $n(l-1) \times n(l-1)$ block gives the following two matrices

$$\mathbf{R} = \begin{bmatrix} \mathbf{I}/2 & \mathbf{N}_\ell \cdots \mathbf{N}_2 \mathbf{N}_1 \\ \mathbf{0}^{n \times n} & \mathbf{I}/2 \end{bmatrix} \quad \text{and} \quad \tilde{\mathbf{R}} = \begin{bmatrix} \mathbf{I}/2 & \tilde{\mathbf{N}}_\ell \cdots \tilde{\mathbf{N}}_2 \tilde{\mathbf{N}}_1 \\ \mathbf{0}^{n \times n} & \mathbf{I}/2 \end{bmatrix}.$$

And we obtain by Theorem A.2 that $\tilde{\mathbf{R}}$ is an $\epsilon/2 + O(\epsilon^2)$ -approximation of \mathbf{R} with respect to $\mathbf{S}_{\mathbf{R}}$. Hence, by Item 3 of Lemma 3.7, we have

$$\tilde{\mathbf{N}}_\ell \cdots \tilde{\mathbf{N}}_2 \tilde{\mathbf{N}}_1 \stackrel{svn}{\approx}_{\epsilon + O(\epsilon^2)} \mathbf{N}_\ell \cdots \mathbf{N}_2 \mathbf{N}_1. \quad \square$$

Lemma 3.17. *If $\tilde{\mathbf{A}} \stackrel{sv}{\approx}_\epsilon \mathbf{A}$, then $\tilde{\mathbf{A}}\tilde{\mathbf{I}} = \mathbf{A}\tilde{\mathbf{I}}$ and $\tilde{\mathbf{A}}^\top \tilde{\mathbf{I}} = \mathbf{A}^\top \tilde{\mathbf{I}}$.*

Proof. Let $x \leftarrow \delta x$ be an arbitrary test vector and let $y = \tilde{\mathbf{I}}$. Then the approximation condition implies

$$\begin{aligned} \left| \delta x^* (\tilde{\mathbf{A}} - \mathbf{A}) \tilde{\mathbf{I}} \right| &\leq \frac{\epsilon}{4} \left(\delta^2 x^* \mathbf{D}_{\text{out}} x - \delta^2 x^* \mathbf{A} \mathbf{D}_{\text{in}}^+ \mathbf{A}^* x + \tilde{\mathbf{I}}^\top \mathbf{D}_{\text{in}} \tilde{\mathbf{I}} - \tilde{\mathbf{I}}^\top \mathbf{A}^\top \mathbf{D}_{\text{out}}^+ \mathbf{A} \tilde{\mathbf{I}} \right) \\ &= \frac{\epsilon}{4} \left(\delta^2 x^* \mathbf{D}_{\text{out}} x - \delta^2 x^* \mathbf{A} \mathbf{D}_{\text{in}}^+ \mathbf{A}^* x \right) \end{aligned}$$

and so taking $\delta \rightarrow 0$ we obtain that $\mathbf{A}\tilde{\mathbf{I}} = \tilde{\mathbf{A}}\tilde{\mathbf{I}}$. An analogous argument setting $x = \tilde{\mathbf{I}}$ shows that $\tilde{\mathbf{I}}^\top \mathbf{A} = \tilde{\mathbf{I}}^\top \tilde{\mathbf{A}}$. \square

Lemma 3.18. *If $\tilde{\mathbf{A}}_i \stackrel{(\mathbf{D}_{\text{in}})_i, (\mathbf{D}_{\text{out}})_i}{\approx}_\epsilon \mathbf{A}_i$ for all $i \in [k]$, letting $\mathbf{D}_{\text{in}} \stackrel{\text{def}}{=} \sum_{i \in [k]} (\mathbf{D}_{\text{in}})_i$ and $\mathbf{D}_{\text{out}} \stackrel{\text{def}}{=} \sum_{i \in [k]} (\mathbf{D}_{\text{out}})_i$, then $\sum_{i \in [k]} \tilde{\mathbf{A}}_i \stackrel{\mathbf{D}_{\text{in}}, \mathbf{D}_{\text{out}}}{\approx}_\epsilon \sum_{i \in [k]} \mathbf{A}_i$. Consequently, if $\tilde{\mathbf{A}}_i \stackrel{sv}{\approx}_\epsilon \mathbf{A}_i$ for every i , then $\sum_{i \in [k]} \tilde{\mathbf{A}}_i \stackrel{sv}{\approx}_\epsilon \sum_{i \in [k]} \mathbf{A}_i$.*

Proof. This follows immediately from Item 3 of Lemma 3.7 being linear over \mathbf{A} and \mathbf{D} . The “consequently” claim follows immediately from noting

$$\left(\sum_{i \in [k]} \mathbf{A} \right) \tilde{\mathbf{I}} = \sum_{i \in [k]} (\mathbf{A}\tilde{\mathbf{I}}) \quad \text{and} \quad \left(\sum_{i \in [k]} \mathbf{A}^\top \right) \tilde{\mathbf{I}} = \sum_{i \in [k]} (\mathbf{A}^\top \tilde{\mathbf{I}}). \quad \square$$

Lemma 3.19. *If $\mathbf{A}_3 \stackrel{\mathbf{D}_{\text{in}}, \mathbf{D}_{\text{out}}}{\approx}_\delta \mathbf{A}_2$ and $\mathbf{A}_2 \stackrel{\mathbf{D}_{\text{in}}, \mathbf{D}_{\text{out}}}{\approx}_\epsilon \mathbf{A}_1$ then $\mathbf{A}_3 \stackrel{\mathbf{D}_{\text{in}}, \mathbf{D}_{\text{out}}}{\approx}_{\epsilon + \delta + \epsilon\delta} \mathbf{A}_1$. Consequently, if $\mathbf{A}_3 \stackrel{sv}{\approx}_\delta \mathbf{A}_2$ and $\mathbf{A}_2 \stackrel{sv}{\approx}_\epsilon \mathbf{A}_1$ then $\mathbf{A}_3 \stackrel{sv}{\approx}_{\epsilon + \delta + \epsilon\delta} \mathbf{A}_1$, and if $\mathbf{A}_3 \stackrel{svn}{\approx}_\delta \mathbf{A}_2$ and $\mathbf{A}_2 \stackrel{svn}{\approx}_\epsilon \mathbf{A}_1$ then $\mathbf{A}_3 \stackrel{svn}{\approx}_{\epsilon + \delta + \epsilon\delta} \mathbf{A}_1$. Moreover, if for $\delta \in (0, 1/2)$ and $\mathbf{A}_0, \dots, \mathbf{A}_\ell$ we have $\mathbf{A}_i \stackrel{sv}{\approx}_{\delta/2\ell} \mathbf{A}_{i-1}$ for every i , then $\mathbf{A}_\ell \stackrel{sv}{\approx}_\delta \mathbf{A}_0$. Moreover, the equivalent claim holds for normalized SV approximation.*

Proof. By Item 3 of Lemma 3.7 we have

$$\begin{bmatrix} \mathbf{D}_{\text{in}} & \mathbf{A}_2^* \\ \mathbf{A}_2 & \mathbf{D}_{\text{out}} \end{bmatrix} \preceq (1 + \varepsilon) \begin{bmatrix} \mathbf{D}_{\text{in}} & \mathbf{A}_1^* \\ \mathbf{A}_1 & \mathbf{D}_{\text{out}} \end{bmatrix}.$$

Thus for arbitrary $x \in \mathbb{C}^m, y \in \mathbb{C}^n$ we have

$$\begin{aligned} & \left| x^* \left(\begin{bmatrix} \mathbf{0} & \mathbf{A}_3 \\ \mathbf{0} & \mathbf{0} \end{bmatrix} - \begin{bmatrix} \mathbf{0} & \mathbf{A}_1 \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \right) y \right| \\ & \leq \left| x^* \left(\begin{bmatrix} \mathbf{0} & \mathbf{A}_3 \\ \mathbf{0} & \mathbf{0} \end{bmatrix} - \begin{bmatrix} \mathbf{0} & \mathbf{A}_2 \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \right) y \right| + \left| x^* \left(\begin{bmatrix} \mathbf{0} & \mathbf{A}_2 \\ \mathbf{0} & \mathbf{0} \end{bmatrix} - \begin{bmatrix} \mathbf{0} & \mathbf{A}_1 \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \right) y \right| \\ & \leq \frac{\delta}{4} \left(x^* \begin{bmatrix} \mathbf{D}_{\text{in}} & \mathbf{A}_2^* \\ \mathbf{A}_2 & \mathbf{D}_{\text{out}} \end{bmatrix} x + y^* \begin{bmatrix} \mathbf{D}_{\text{in}} & \mathbf{A}_2^* \\ \mathbf{A}_2 & \mathbf{D}_{\text{out}} \end{bmatrix} y \right) + \frac{\varepsilon}{4} \left(x^* \begin{bmatrix} \mathbf{D}_{\text{in}} & \mathbf{A}_1^* \\ \mathbf{A}_1 & \mathbf{D}_{\text{out}} \end{bmatrix} x + y^* \begin{bmatrix} \mathbf{D}_{\text{in}} & \mathbf{A}_1^* \\ \mathbf{A}_1 & \mathbf{D}_{\text{out}} \end{bmatrix} y \right) \\ & \leq \frac{\delta(1 + \varepsilon)}{4} \left(x^* \begin{bmatrix} \mathbf{D}_{\text{in}} & \mathbf{A}_1^* \\ \mathbf{A}_1 & \mathbf{D}_{\text{out}} \end{bmatrix} x + y^* \begin{bmatrix} \mathbf{D}_{\text{in}} & \mathbf{A}_1^* \\ \mathbf{A}_1 & \mathbf{D}_{\text{out}} \end{bmatrix} y \right) + \frac{\varepsilon}{4} \left(x^* \begin{bmatrix} \mathbf{D}_{\text{in}} & \mathbf{A}_1^* \\ \mathbf{A}_1 & \mathbf{D}_{\text{out}} \end{bmatrix} x + y^* \begin{bmatrix} \mathbf{D}_{\text{in}} & \mathbf{A}_1^* \\ \mathbf{A}_1 & \mathbf{D}_{\text{out}} \end{bmatrix} y \right) \\ & = \frac{\delta + \varepsilon + \varepsilon\delta}{4} \left(x^* \begin{bmatrix} \mathbf{D}_{\text{in}} & \mathbf{A}_1^* \\ \mathbf{A}_1 & \mathbf{D}_{\text{out}} \end{bmatrix} x + y^* \begin{bmatrix} \mathbf{D}_{\text{in}} & \mathbf{A}_1^* \\ \mathbf{A}_1 & \mathbf{D}_{\text{out}} \end{bmatrix} y \right) \end{aligned}$$

and so we conclude by Item 3 of Lemma 3.7. The first “consequently” claim follows from the fact that by Lemma 3.17 we have $\mathbf{A}_2 \vec{1} = \mathbf{A} \vec{1}$ and $\mathbf{A}_2^\top \vec{1} = \mathbf{A}_1^\top \vec{1}$, and the second follows immediately.

For the “moreover” claim, we prove by induction on i that $\mathbf{A}_i \overset{sv}{\approx}_{i\delta/2\ell + i\delta^2/\ell} \mathbf{A}_0$, which suffices as $\ell\delta/2\ell + \ell\delta^2/\ell \leq \delta$. Assuming this holds for i , we have

$$\mathbf{A}_{i+1} \overset{sv}{\approx}_{\delta/2\ell} \mathbf{A}_i \overset{sv}{\approx}_{i\delta/2\ell + i\delta^2/\ell} \mathbf{A}_0$$

and thus we have

$$\mathbf{A}_{i+1} \overset{sv}{\approx}_{(i+1)\delta/2\ell + \gamma} \mathbf{A}_0$$

for

$$\gamma = \frac{i\delta^2}{\ell} + \frac{\delta}{2\ell} \left(\frac{i\delta}{2\ell} + \frac{i\delta^2}{\ell} \right) \leq \frac{i\delta^2}{\ell} + \delta^2/2\ell + \delta^3/2\ell^2 \leq \frac{(i+1)\delta^2}{\ell}$$

as desired. The proof for normalized SV approximation is essentially identical, so we omit it. \square

Lemma 3.20. *Let G be a strongly connected, d -regular directed multigraph on n vertices with adjacency matrix \mathbf{A} and let $\mathbf{J} \in \mathbb{R}^{n \times n}$ be a matrix with $1/n$ in every entry (i.e. \mathbf{J} is the walk matrix of the complete graph with a self loop on every vertex). Then $\lambda(G) \leq 1 - \lambda/2$ if and only if $\mathbf{A}/d \overset{svn}{\approx}_\lambda \mathbf{J}$.*

Proof. Note that by definition of $\mathbf{A} \overset{svn}{\approx}_\lambda \mathbf{J}$,

$$\begin{aligned} |x^*(\mathbf{A}/d - \mathbf{J})y| & \leq \frac{\lambda}{4} (x^*(\mathbf{I} - \mathbf{J}\mathbf{J}^*)x + y^*(\mathbf{I} - \mathbf{J}^*\mathbf{J})y) \\ & = \frac{\lambda}{4} (x^*(\mathbf{I} - \mathbf{J})x + y^*(\mathbf{I} - \mathbf{J})y) \end{aligned}$$

Therefore, $\mathbf{A}/d \overset{svn}{\approx}_\lambda \mathbf{J}$ if and only if $\mathbf{A}/d \overset{\circ}{\approx}_{\lambda/2} \mathbf{J}$. By [AKM⁺20, Lemma 5.2] $\mathbf{A}/d \overset{\circ}{\approx}_{\lambda/2} \mathbf{J}$ if and only if $\lambda(G) \leq 1 - \lambda/2$, so the result follows. \square

We show a slightly stronger version of Item 2 of Lemma 3.7 for non-negative matrices, in that we show SV sparsification of the bipartite lift of a graph implies SV sparsification of the underlying directed graph, even if the sparsifier is not promised to be bipartite-preserving, and even if the sparsifier is not itself undirected.

Lemma A.5. *Given non-negative $\mathbf{A} \in \mathbb{R}_{\geq 0}^{m \times n}$, suppose there is nonnegative $\tilde{\mathbf{A}} = \begin{bmatrix} \tilde{\mathbf{A}}_{F,F} & \tilde{\mathbf{A}}_{F,F^c} \\ \tilde{\mathbf{A}}_{F^c,F} & \tilde{\mathbf{A}}_{F^c,F^c} \end{bmatrix}$*

such that $\tilde{\mathbf{A}} \overset{sv}{\approx}_\varepsilon \begin{bmatrix} \mathbf{0} & \mathbf{A}^\top \\ \mathbf{A} & \mathbf{0} \end{bmatrix}$. Then

1. $\tilde{\mathbf{A}}_{F,F} = \mathbf{0}$ and $\tilde{\mathbf{A}}_{F^c,F^c} = \mathbf{0}$.
2. $\tilde{\mathbf{A}}_{F^c,F} \overset{sv}{\approx}_\varepsilon \mathbf{A}$ and $\tilde{\mathbf{A}}_{F,F^c} \overset{sv}{\approx}_\varepsilon \mathbf{A}^\top$.

Proof. Let $\mathbf{D}_{\text{out}} = \text{diag}(\mathbf{A}\vec{\mathbf{1}})$ and $\mathbf{D}_{\text{in}} = \text{diag}(\vec{\mathbf{1}}^\top \mathbf{A})$ and observe $\vec{\mathbf{1}}^\top \mathbf{D}_{\text{out}} \vec{\mathbf{1}} - \vec{\mathbf{1}}^\top \mathbf{A} \mathbf{D}_{\text{in}}^+ \mathbf{A} \vec{\mathbf{1}} = \vec{\mathbf{1}}^\top \mathbf{D}_{\text{in}} \vec{\mathbf{1}} - \vec{\mathbf{1}}^\top \mathbf{A}^\top \mathbf{D}_{\text{out}}^+ \mathbf{A} \vec{\mathbf{1}} = 0$.

1. Let $x \in \mathbb{C}^n$ be arbitrary, and consider the test vectors $x' = \delta \begin{bmatrix} x \\ 0^m \end{bmatrix}$, $y' = \begin{bmatrix} \vec{\mathbf{1}} \\ \vec{\mathbf{1}} \end{bmatrix}$. Then from the definition of SV approximation we obtain

$$\begin{aligned} \left| \delta x^\top \tilde{\mathbf{A}}_{F,F} \vec{\mathbf{1}} + \delta x^* \tilde{\mathbf{A}}_{F,F^c} \vec{\mathbf{1}} - \delta x \mathbf{A}^\top \vec{\mathbf{1}} \right| &= \left| x'^* \left(\begin{bmatrix} \mathbf{0} & \mathbf{A}^\top \\ \mathbf{A} & \mathbf{0} \end{bmatrix} - \tilde{\mathbf{A}} \right) y' \right| \\ &\leq \frac{\varepsilon}{2} \cdot \delta^2 x^* (\mathbf{D}_{\text{out}} - \mathbf{A} \mathbf{D}_{\text{in}}^+ \mathbf{A}) x \end{aligned}$$

and hence $\tilde{\mathbf{A}}_{F,F} \vec{\mathbf{1}} + x \tilde{\mathbf{A}}_{F,F^c} \vec{\mathbf{1}} = x \mathbf{A}^\top \vec{\mathbf{1}}$ by taking $\delta \rightarrow 0$. Considering the test vectors $x' = \delta \begin{bmatrix} x \\ 0^m \end{bmatrix}$, $y'' = \begin{bmatrix} \vec{\mathbf{1}} \\ -\vec{\mathbf{1}} \end{bmatrix}$ we likewise obtain $x \tilde{\mathbf{A}}_{F,F} \vec{\mathbf{1}} - x \tilde{\mathbf{A}}_{F,F^c} \vec{\mathbf{1}} = -x \mathbf{A}^\top \vec{\mathbf{1}}$ and hence $x \tilde{\mathbf{A}}_{F,F} \vec{\mathbf{1}} = 0$.

By an analogous argument we obtain $\tilde{\mathbf{A}}_{F^c,F^c} = 0$.

2. For arbitrary $x \in \mathbb{C}^m$, $y \in \mathbb{C}^n$, apply the test vectors $[0^n, x]$, $[y, 0^m]$. Then we obtain

$$\left| x^* (\mathbf{A} - \tilde{\mathbf{A}}_{F^c,F}) y \right| \leq \frac{\varepsilon}{2} (x^* (\mathbf{D} - \mathbf{A} \mathbf{D}^+ \mathbf{A}^T) x + y^* (\mathbf{D} - \mathbf{A}^T \mathbf{D}^+ \mathbf{A}) y)$$

thus $\tilde{\mathbf{A}}_{F^c,F} \overset{sv}{\approx}_\varepsilon \mathbf{A}$ and an analogous argument shows $\tilde{\mathbf{A}}_{F,F^c} \overset{sv}{\approx}_\varepsilon \mathbf{A}^\top$.

□

A.3 Separations

Proposition 3.12. *For every $\alpha, \varepsilon \in (0, 1)$, there are symmetric matrices \mathbf{W} , $\tilde{\mathbf{W}}$ with $\|\mathbf{W}\|, \|\tilde{\mathbf{W}}\| \leq 1$ such that $\tilde{\mathbf{W}} \overset{\circ}{\approx}_\varepsilon \mathbf{W}$ but $\tilde{\mathbf{W}}$ is not an ε SV-approximation of \mathbf{W} with respect to $\mathbf{I} - \mathbf{W}$ for any $\varepsilon' < \frac{\varepsilon}{\sqrt{1-\alpha^2}}$.*

Proof. Given α , we define

$$\mathbf{W} = \begin{bmatrix} \alpha & 0 \\ 0 & -\alpha \end{bmatrix}$$

and

$$\tilde{\mathbf{W}} = \begin{bmatrix} \alpha & \varepsilon \sqrt{1-\alpha^2} \\ \varepsilon \sqrt{1-\alpha^2} & -\alpha \end{bmatrix}.$$

We first show that $\widetilde{\mathbf{W}} \overset{\circ}{\approx}_\varepsilon \mathbf{W}$. Since both matrices are symmetric, suffices [AKM⁺20, Lemma 3.7] to show the equivalent statement for PSD approximation between $\mathbf{I} - \widetilde{\mathbf{W}}, \mathbf{I} - \mathbf{W}$ and $\mathbf{I} + \widetilde{\mathbf{W}}, \mathbf{I} + \mathbf{W}$, i.e.

$$(1 - \varepsilon)(\mathbf{I} - \mathbf{W}) \preceq \mathbf{I} - \widetilde{\mathbf{W}} \preceq (1 + \varepsilon)(\mathbf{I} - \mathbf{W})$$

$$(1 - \varepsilon)(\mathbf{I} + \mathbf{W}) \preceq \mathbf{I} + \widetilde{\mathbf{W}} \preceq (1 + \varepsilon)(\mathbf{I} + \mathbf{W}).$$

All four inequalities are implied by

$$\begin{bmatrix} \varepsilon(1 - \alpha) & \varepsilon\sqrt{1 - \alpha^2} \\ \varepsilon\sqrt{1 - \alpha^2} & \varepsilon(1 + \alpha) \end{bmatrix} \succeq \mathbf{0}.$$

Using Fact A.1 this is equivalent to $\varepsilon(1 - \alpha) - \varepsilon(1 - \alpha^2)(1 + \alpha)^{-1} \geq 0$ which holds with equality, so $\widetilde{\mathbf{W}} \overset{\circ}{\approx}_\varepsilon \mathbf{W}$. Now assume that $\widetilde{\mathbf{W}} \overset{svn}{\approx}_{\varepsilon'} \mathbf{W}$ for some ε' . From the definition of SV approximation applied to test vector $z = \begin{bmatrix} x \\ x \end{bmatrix}$, we have

$$\pm(\widetilde{\mathbf{W}} - \mathbf{W}) \preceq \varepsilon' (\mathbf{I} - \mathbf{W}^2) = \varepsilon'(1 - \alpha^2)\mathbf{I}.$$

This is equivalent to

$$\varepsilon\sqrt{1 - \alpha^2} \leq \varepsilon'(1 - \alpha^2).$$

Thus for $\widetilde{\mathbf{W}} \overset{svn}{\approx}_\varepsilon \mathbf{W}$, a necessary condition is for $\varepsilon' \geq \frac{\varepsilon}{\sqrt{1 - \alpha^2}}$. \square

B Facts about Singular Values

Lemma B.1. *Let $\sigma_i(\cdot)$ denote the i th largest singular value. For any matrix $\mathbf{A} \in \mathbb{C}^{n \times n}$ and any positive integer k ,*

$$\sigma_2(\mathbf{A}^k) \leq \sigma_2(\mathbf{A}) \cdot \sigma_1(\mathbf{A})^{k-1}.$$

Proof. By the variational characterization of singular values, we have

$$\sigma_2(\mathbf{A}^k) = \min_{v \in \mathbb{C}^n} \max_{x \perp v} \frac{\|\mathbf{A}^k x\|}{\|x\|} \leq \min_{v \in \mathbb{C}^n} \max_{x \perp v} \frac{\|\mathbf{A}^{k-1}\| \cdot \|\mathbf{A}x\|}{\|x\|} \leq \|\mathbf{A}\|^{k-1} \cdot \min_{v \in \mathbb{C}^n} \max_{x \perp v} \frac{\|\mathbf{A}x\|}{\|x\|} = \sigma_2(\mathbf{A}) \cdot \sigma_1(\mathbf{A})^{k-1}.$$

\square

Lemma B.2. *Let $\sigma_i(\cdot)$ denote the i th largest singular value. For any matrix $\mathbf{A}, \mathbf{B} \in \mathbb{C}^{n \times n}$,*

$$\sigma_2(\mathbf{A} + \mathbf{B}) \leq \sigma_2(\mathbf{A}) + \sigma_1(\mathbf{B}).$$

Furthermore, if \mathbf{A}, \mathbf{B} share a common right singular vector that achieves the maximum singular value in each of them respectively, then

$$\sigma_2(\mathbf{A} + \mathbf{B}) \leq \sigma_2(\mathbf{A}) + \sigma_2(\mathbf{B})$$

Proof. For the first part of the statement, by the variational characterization of singular values, we have

$$\sigma_2(\mathbf{A} + \mathbf{B}) = \min_{v \in \mathbb{C}^n} \max_{x \perp v} \frac{\|(\mathbf{A} + \mathbf{B})x\|}{\|x\|} \leq \min_{v \in \mathbb{C}^n} \max_{x \perp v} \frac{\|\mathbf{A}x\| + \|\mathbf{B}x\|}{\|x\|} \leq \min_{v \in \mathbb{C}^n} \max_{x \perp v} \frac{\|\mathbf{A}x\|}{\|x\|} + \|\mathbf{B}\|.$$

For the second part of the statement, let v denote a common right singular vector of \mathbf{A}, \mathbf{B} that respectively achieves the maximum singular value in each of them. Then we have

$$\begin{aligned}\sigma_2(\mathbf{A} + \mathbf{B}) &= \min_{v' \in \mathbb{C}^n} \max_{x \perp v'} \frac{\|(\mathbf{A} + \mathbf{B})x\|}{\|x\|} \leq \max_{x \perp v} \frac{\|(\mathbf{A} + \mathbf{B})x\|}{\|x\|} \\ &\leq \max_{x \perp v} \frac{\|\mathbf{A}x\| + \|\mathbf{B}x\|}{\|x\|} \leq \max_{x \perp v} \frac{\|\mathbf{A}x\|}{\|x\|} + \max_{x \perp v} \frac{\|\mathbf{B}x\|}{\|x\|} = \sigma_2(\mathbf{A}) + \sigma_2(\mathbf{B}).\end{aligned}$$

The third part of the statement has essentially the same proof as the second. \square

We now use these properties to show that small perturbations preserve the smallest singular value:

Lemma B.3. *Let $\sigma_2(\cdot)$ denote the second largest singular value. Let G be an Eulerian graph such that $\sigma_2(\mathbf{D}_G^{-1/2} \mathbf{A}_G \mathbf{D}_G^{-1/2}) \leq 1 - 1/\gamma$. For every H such that $\mathbf{D}_H = \mathbf{D}_G$ and $\|\mathbf{A}_H - \mathbf{A}_G\| \leq \delta$, we have $\sigma_2(\mathbf{D}_H^{-1/2} \mathbf{A}_H \mathbf{D}_H^{-1/2}) \leq 1 - 1/\gamma + \delta/d_{\min}$ where d_{\min} is the minimum diagonal entry of \mathbf{D}_G .*

Proof. By lemma B.2,

$$\begin{aligned}\sigma_2(\mathbf{D}_H^{-1/2} \mathbf{A}_H \mathbf{D}_H^{-1/2}) &\leq \sigma_2(\mathbf{D}_G^{-1/2} \mathbf{A}_G \mathbf{D}_G^{-1/2}) + \|\mathbf{D}_G^{-1/2} (\mathbf{A}_H - \mathbf{A}_G) \mathbf{D}_G^{-1/2}\| \\ &\leq 1 - 1/\gamma + \|\mathbf{D}_G^{-1/2}\|^2 \cdot \|\mathbf{A}_H - \mathbf{A}_G\| \\ &= 1 - 1/\gamma + \delta/d_{\min}.\end{aligned}\quad \square$$

Lemma B.4. *Let $\mathbf{A}, \tilde{\mathbf{A}} \in \mathbb{R}^{n \times n}$ be the adjacency matrices of Eulerian graphs with no isolated vertices. Suppose $\tilde{\mathbf{A}} \stackrel{sv}{\approx}_\varepsilon \mathbf{A}$. Let \mathbf{D} be their diagonal matrix of degrees. Then*

$$1 - \sigma_2(\mathbf{D}^{-1/2} \tilde{\mathbf{A}} \mathbf{D}^{-1/2}) \leq (1 + 2\varepsilon) \cdot (1 - \sigma_2(\mathbf{D}^{-1/2} \mathbf{A} \mathbf{D}^{-1/2})).$$

Proof. Define $\mathbf{N} = \mathbf{D}^{-1/2} \mathbf{A} \mathbf{D}^{-1/2}$ and $\tilde{\mathbf{N}} = \mathbf{D}^{-1/2} \tilde{\mathbf{A}} \mathbf{D}^{-1/2}$.

It suffices to show

$$\pm(\sigma_2(\tilde{\mathbf{N}}) - \sigma_2(\mathbf{N})) \leq \sigma_2(\tilde{\mathbf{N}} - \mathbf{N}) \leq 2\varepsilon \cdot (1 - \sigma_2(\mathbf{N})).$$

The first inequality follows from lemma B.2. For the second inequality, note that the definition of SV approximation can be rewritten as for all $x \in \mathbb{C}^n$,

$$\|(\tilde{\mathbf{N}} - \mathbf{N})x\| \leq \varepsilon \cdot (1 - \|\mathbf{N}x\|^2).$$

Note that $\tilde{\mathbf{N}}, \mathbf{N}$ have a common left and right eigenvector given by $v = \mathbf{D}^{1/2} \vec{1}$, and this achieves their respective maximum singular values of 1.⁷ Hence, if we wish to maximize the LHS over all unit vectors x , it suffices to maximize over all unit vectors $x \perp v$. Doing so, we obtain

$$\|\tilde{\mathbf{N}} - \mathbf{N}\| \leq \varepsilon \cdot (1 - \sigma_2(\mathbf{N})^2) \leq 2\varepsilon \cdot (1 - \sigma_2(\mathbf{N}))$$

where the last inequality is because $\|\mathbf{N}\| \leq 1$ ([CKP⁺17] lemma B.4). \square

⁷We know it corresponds to a singular value of 1 and this its not possible to have a singular value larger than 1 for this matrix ([CKP⁺17] lemma B.4).

Lemma B.5. Let $\sigma_i(\cdot)$ denote the i th largest singular value. Let $\mathbf{A} \in \mathbb{R}^{n \times n}$ be the adjacency matrix of a strongly connected Eulerian graph G with diagonal degree matrix \mathbf{D} and no isolated vertices. Let $\mathbf{N} = \mathbf{D}^{-1/2} \mathbf{A} \mathbf{D}^{-1/2}$. Let λ denote the second largest eigenvalue of $\mathbf{S}_{\mathbf{N}}$.⁸ Then we have

$$\sigma_2((1 - \gamma)\mathbf{N} + \gamma\mathbf{I}) \leq 1 - (1 - \lambda)\gamma + O(\gamma^2).$$

In particular, if $\gamma \geq 1/\text{poly}(n)$ and \mathbf{A} has $\text{poly}(n)$ weak mixing time and all weights are specified using $O(\log n)$ bits of precision, then

$$\sigma_2((1 - \gamma)\mathbf{N} + \gamma\mathbf{I}) \leq 1 - 1/\text{poly}(n).$$

Proof. Let v denote a singular vector of $(1 - \gamma)\mathbf{N} + \gamma\mathbf{I}$ corresponding to its second-largest singular value. We then have

$$\begin{aligned} [\sigma_2((1 - \gamma)\mathbf{N} + \gamma\mathbf{I})]^2 &= (1 - \gamma)^2 v^* \mathbf{N}^* \mathbf{N} v + 2\gamma(1 - \gamma) v^* \mathbf{S}_{\mathbf{N}} v + \gamma^2 \\ &\leq (1 - \gamma)^2 + 2\gamma(1 - \gamma)\lambda + \gamma^2 \\ &\leq 1 - 2(1 - \lambda)\gamma + O(\gamma^2). \end{aligned}$$

In the above, we used the fact that $\|\mathbf{N}\| \leq 1$ ([CKP⁺17] lemma B.4).

To show the second part of the statement, let $\tau(\mathbf{M})$ denote the weak mixing time of the random walk on the graph corresponding to any Eulerian adjacency matrix \mathbf{M} . Define $\mathbf{L} = \mathbf{I} - \mathbf{N}$. Note that \mathbf{L} is related to $\mathbf{I} - \mathbf{A} \mathbf{D}^{-1}$ by a change of basis with polynomial condition number. We may assume the graph is strongly connected as this is necessary for the weak mixing time to be finite. It suffices to prove

$$\frac{1}{1 - \lambda} \leq \text{poly}(n) \cdot \tau(\mathbf{N}).$$

We do so by showing

$$\frac{1}{1 - \lambda} \leq \text{poly}(n) \cdot \tau(\mathbf{S}_{\mathbf{N}}) \leq \text{poly}(n) \cdot \|\mathbf{S}_{\mathbf{L}}^+\| \leq \text{poly}(n) \cdot \|\mathbf{L}^+\| \leq \text{poly}(n) \cdot \tau(\mathbf{N}).$$

The first inequality is just the folklore result relating spectral gap to weak mixing time with an $O(\log n)$ -factor loss. The second and fourth inequalities are from ([CKP⁺16] Theorem 21).

We now show the third inequality, that $\|\mathbf{S}_{\mathbf{L}}^+\| \leq \text{poly}(n) \cdot \|\mathbf{L}^+\|$. We have

$$\|\mathbf{S}_{\mathbf{L}}^+\| \leq \text{poly}(n) \cdot \|(\mathbf{L}^T \mathbf{S}_{\mathbf{L}}^+ \mathbf{L})^+\| = \text{poly}(n) \cdot \|\mathbf{S}_{\mathbf{L}^+}\| \leq \text{poly}(n) \cdot (\|\mathbf{L}^+\| + \|\mathbf{L}^{T^+}\|)/2 = \text{poly}(n) \cdot \|\mathbf{L}^+\|$$

where the first inequality in the line immediately above is a corollary of ([CKP⁺16] Lemma 13). \square

Lemma B.6. For matrices $\mathbf{A}, \mathbf{B} \in \mathbb{C}^{n \times n}$,

$$\|\mathbf{AB}\| = \|(\mathbf{A}^* \mathbf{A})^{1/2} \mathbf{B}\|, \|\mathbf{BA}\| = \|\mathbf{B}(\mathbf{A} \mathbf{A}^*)^{1/2}\|.$$

Proof. For any matrix \mathbf{M} ,

$$\|\mathbf{M}\| = \sqrt{\lambda_{\max}(\mathbf{M}^* \mathbf{M})} = \max_{x \in \mathbb{C}^m - \{\vec{0}\}} \frac{x^* \mathbf{M}^* \mathbf{M} x}{x^* x}.$$

Thus,

$$\|\mathbf{BA}\| = \max_{x \in \mathbb{C}^m - \{\vec{0}\}} \frac{x^* \mathbf{B}^* \mathbf{A}^* \mathbf{A} \mathbf{B} x}{x^* x} = \max_{x \in \mathbb{C}^m - \{\vec{0}\}} \frac{x^* \mathbf{B}^* (\mathbf{A}^* \mathbf{A})^{1/2} (\mathbf{A}^* \mathbf{A})^{1/2} \mathbf{B} x}{x^* x} = \|(\mathbf{A}^* \mathbf{A})^{1/2} \mathbf{B}\|.$$

The other equality is proved similarly. \square

⁸Here we mean largest according to the actual eigenvalues, not their magnitudes.